



Physician Orders

LEB Severe Sepsis Early Management Plan

[X or R] = will be ordered unless marked out.

PEDIATRIC

Height: _____ cm Weight: _____ kg

Allergies: _____ [] No known allergies

[] Latex allergy [] Other: _____

Patient Care

[] Intermittent Needle Therapy Insert/Site T;N, STAT, q2h(std), If no IV access available
Care LEB (INT Insert/Site Care LEB)

Nursing Communication

[] Nursing Communication T;N, Discontinue NS Bolus PRN order if not given by Sepsis Response Team

Respiratory Care

[] Oxygen Delivery T;N, Routine, Special Instructions: Titrate to keep O2 Sat \geq 92%. Wean to room air.

[] ISTAT POC (RT Collect) T;N, STAT once, Test Select Lactate, CBG

Continuous Infusions

[] Sodium Chloride 0.9% (NS Order Bolus) T;N, 20 mL/kg, Injection, IV, once, STAT, (infuse over 15 min).

[] Sodium Chloride 0.9% (NS Order Bolus) T;N, 20 mL/kg, Injection, IV, PRN, (for 2 doses), STAT, (infuse over 15 min). Confirm need for additional doses with MD/LIP prior to administration.

[] Sodium Chloride 0.9% (NS Order Bolus) T;N, 20 mL/kg, Injection, IV, once, STAT, (infuse over 30 min).

[] Sodium Chloride 0.9% (NS Order Bolus) T;N, 20 mL/kg, Injection, IV, PRN, (for 2 doses), STAT, (infuse over 30 min). Confirm need for additional doses with MD/LIP prior to administration.

Medications

NOTE: Empiric Treatment for Bloodstream Infections (reassess within 48 hours)

[] vancomycin _____ mg (15 mg/kg), Injection, Ped Injectable, IVPB, once, STAT, T;N

[] vancomycin _____ mg (15 mg/kg), Injection, Ped Injectable, IVPB, q6h (for 14 day),
Routine, T;N

NOTE: LESS THAN 5 days hospitalization AND low risk for Multi-Drug Resistant Organisms

[] cefotaxime _____ mg (50 mg/kg), Injection, IV, once, STAT, T;N

[] cefotaxime _____ mg (50 mg/kg), Injection, IV, q8h (for 14 day), Routine, T;N

[] cefTRIAXone _____ mg (100 mg/kg), Injection, IV, once, STAT, T;N

[] cefTRIAXone _____ mg (100 mg/kg), Injection, IV, q24h (for 14 day), Routine, T;N

NOTE: GREATER THAN OR EQUAL TO 5 days hospitalization OR high risk for Multi-Drug Resistant Organisms

[] ceftazidime _____ mg (50 mg/kg), Injection, IV, once, STAT, T;N

[] ceftazidime _____ mg (50 mg/kg), Injection, IV, q8h (for 14 day), Routine, T;N

NOTE: Suspected intra-abdominal focus of infection

[] meropenem _____ mg (20 mg/kg), Injection, IV, once, STAT, T;N

[] meropenem _____ mg (20 mg/kg), Injection, IV, q8h (for 14 day), Routine, T;N



attach patient label



Physician Orders

LEB Severe Sepsis Early Management Plan

[X or R] = will be ordered unless marked out.

PEDIATRIC

Laboratory		
<input type="checkbox"/>	Blood Culture	T;N, STAT, once, Specimen Source: Peripheral Blood
<input type="checkbox"/>	Blood Culture	T;N, STAT, once, Specimen Source: Line, Central
<input type="checkbox"/>	CBC with diff	T;N, STAT, Type: Blood
<input type="checkbox"/>	CRP	T;N, STAT, Type: Blood
Consults		
<input type="checkbox"/>	Consult MD Group	T;N, Routine, Group: ULPS Critical Care, Reason for Consult: Suspected sepsis

Date

Time

Physician's Signature

MD Number

LEB Cross Severe Sepsis Early Mngt-42225-PP-
QM122112-043013