

Physician Orders

LEB Opthamology Pre Op Plan

[X or R] = will be ordered unless marked out.

	PEDIATRIC					
Heigh	t:cm	kg				
Allergies:		[] No known allergies				
		• • • •				
[X]	Initiate Powerplan Phase	T;N, Phase: LEB Ophthamology	/ Preop Phase			
	· ·	Admission/Transfer/Discharg	ge			
[]	Patient Status Initial Inpatient	Attending Physician:				
	Bed Type: [] Med Surg []Critical Care	[] Stepdown [] Other				
[]	Patient Status Initial Outpatient	Attending Physician:				
	Outpatient Status/Service: [] OP-A[] OP-Diagnostic Procedure [] OP-Observation Services					
	Initial status – inpatient For a condition/dx with severity of illness or co-morbid conditions indicating a hospital stay greater than 24 hours					
	is required.					
	Initial Status Outpatient – Ambulatory surgery – Outpatient surgery/procedure with discharge anticipated after a routine or, in some cases,					
	extended recovery. • Routine recovery after outpatient surgery is estimated at 6-8 hours.					
	• "Extended" routine recovery may be required for a patient to stay longer (could be overnight) to recover from anticipated sequela of					
	surgery including effects of anesthesia, nausea, pain.					
	• For unanticipated sequela of surgery or a complicated post operative course, the patient may require a status change to inpatient. Please					
	consult with a case manager before making this choice of "status change".					
	• Examples: Initial status outpatient is generally selected for patients undergoing PCI, diagnostic caths, EP studies, ablations, pacemaker implantations, other routine surgeries.					
	Initial status Outpatient -Observation Services – Short term treatment, assessment and reassessment - estimate discharge within 24 hours					
	 In some cases (for Medicare patients), this can be extended to 48 hours. Observation Services can also be utilized when it is unclear (without additional assessment) whether the patient will require an inpatient 					
	stay.					
[]	Notify Physician Once	T;N, of room number on arrival	to unit			
[]	Vital Signs	T;N, Routine Monitor and Recor	d T,P,R,BP, per routine			
	•	Activity				
[]	Out Of Bed (Activity As Tolerated)	T;N, Up Ad Lib				
		Food/Nutrition				
[]	NPO	Start at: T;N				
	Patient Care					
[]	Isolation Precautions	T;N, per policy				
[]	O2 Sat Spot Check-NSG	T;N, with vital signs				
[]	Cardiopulmonary Monitor	T;N Routine, Monitor Type: CP	Monitor			
Medications						
[]	proparacaine ophthalmic 0.5 % solution	1 drop, Ophthalmic Soln,	eye (s), once, STAT, T;N			
[]	phenylephrine ophthalmic 2.5% solution	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			
[]	tropicamide 1% ophthalmic solution	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			
[]	cyclopentolate ophthalmic 1% solution	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			
[]	cyclopentolate-phenylephrine opthalmic	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			
[]	flurbiprofen 0.03% ophthalmic solution	1 drop, Ophthalmic Soln, on arrival. STAT,T;N	eye (s),on call to OR, 1 drop to be administered			
L 1	ciproflovacia 0.3% on the latic colution	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			
	ciprofloxacin 0.3% ophthalmic solution					
[]	moxifloxacin 0.5% ophthalmic solution	1 drop, Ophthalmic Soln,	_ eye(s), q5min, (3 doses), STAT, T;N			





Physician Orders

LEB Opthamology Pre Op Plan

[X or R] = will be ordered unless marked out.

Laboratory					
[]	Hematocrit (Hct)	STAT, T;N, once, Type: Blood			
[]	Pregnancy Screen Serum	STAT, T;N, Type: Blood			
Consults/Notifications					

Date

Time

Physician's Signature

MD Number

LEB OPTHA Pre-Op-42601 PP-QM1008 091312