



Physician Orders ADULT: Adult Procedural Sedation Plan

Adult Procedural Sedation Plan

Non Categorized

R Procedural Moderate Sedation

Patient Care

- ☒ IV Insert/Site Care
Routine

Nursing Communication

- ☒ Nursing Communication
Adult Procedural Sedation: If reversal agent administered, please verify that event is entered into the sedation database.(Procedure Order)
- ☒ Nursing Communication
Adult Procedural Sedation: Discontinue Adult Procedural Sedation Plan orders when patient discharged from Procedural Area. (Procedure Order)

Continuous Infusion

- ☐ Sodium Chloride 0.9%
1,000 mL, IV, Routine, 75 mL/hr
- ☐ Sodium Chloride 0.9%
1,000 mL, IV, Routine, Titrate, TKO, Start at 20 mL/hr

Medications

Deep Sedating Agents - Deep Sedation Privileged MDs ONLY(NOTE)*

- ☐ ketamine
1 mg/kg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY
- ☐ etomidate
0.1 mg/kg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY
- ☐ propofol bolus (for procedural anesthesia)
1 mg/kg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY
- ☐ propofol bolus (for procedural anesthesia)
0.5 mg/kg, Injection, IV Push, q5min, PRN Sedation, STAT, (for 2 dose), To Follow 1 mg/kg initial dose as needed. In procedure ONLY.
- Moderate Sedating Agents - Moderate Sedation Privileged MDs ONLY(NOTE)*
NOTE: Consider lower doses for patients greater than 65 years old or opiate naive.(NOTE)*
- ☐ fentaNYL
50 mcg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY (MAX initial dose = 50 mcg)
- ☐ fentaNYL
50 mcg, Injection, IV Push, q5min, PRN Sedation, STAT, (for 3 dose), For Sedation in procedure ONLY
Comments: Sedation in procedure ONLY. (Max initial dose = 50 mcg),(Max dose = 500 mcg)
- ☐ midazolam
1 mg, Injection, IV Push, N/A, STAT, For sedation in procedure ONLY. (Max dose = 5 mg)
- ☐ midazolam
2 mg, Injection, IV Push, N/A, STAT, For sedation in procedure ONLY. (Max dose = 5 mg)
- ☐ midazolam
1 mg, Injection, IV Push, q5min, PRN Sedation, STAT, (for 4 dose), For sedation in procedure ONLY (Max dose = 5 mg)
- ☐ morphine
4 mg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY. (Max dose = 10 mg)
- ☐ morphine
2 mg, Injection, IV Push, q5min, PRN Sedation, STAT, (for 5 dose), For sedation in procedure ONLY. (Max dose = 10 mg)
- ☐ HYDROmorphine





Physician Orders ADULT: Adult Procedural Sedation Plan

1 mg, Injection, IV Push, N/A, Sedation, STAT, For Sedation in procedure ONLY

Comments: For sedation in procedure ONLY. (Max dose = 4 mg) Note: HYDROmorphone 1 mg = morPHINE 8 mg.

- ☐ HYDROmorphone
0.5 mg, Injection, IV Push, q5min, PRN Sedation, STAT, (for 2 dose), For Sedation in procedure ONLY
Comments: For sedation in procedure ONLY. Max dose = 4 mg. Note: HYDROmorphone 0.5 mg = morPHINE 4 mg
- Reversal Agents(NOTE)*
NOTE: If reversal agent administered, please enter event into sedation database(NOTE)*
- ☐ naloxone
0.2 mg, Injection, IV Push, q1min, PRN Other, specify in Comment, STAT, (for 4 dose)
Comments: Reversal agent for opioid oversedation. If used, document in sedation database. For Procedure Use Only
- ☐ flumazenil
0.2 mg, Injection, IV Push, q1min, PRN Other, specify in Comment, STAT, (for 4 dose)
Comments: Reversal agent for benzodiazepine oversedation/respiratory depression If used, document in sedation database. For Procedure use ONLY

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

*Report Legend:

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

