



Physician Orders ADULT: OB Fetal Demise Plan

Initiate Orders Phase

Non Categorized

- ☒ Initiate Powerplan Phase

Phase: Fetal Demise (Cytotec) Phase, When to Initiate: _____

OB Fetal Demise Phase

Activity

- ☒ Bedrest

maintain patient in the lateral recumbant position with head slightly elevated for 2 hours after insertion of misoprostol (Cytotec).

- ☒ Bedrest w/BRP

prior to insertion of misoprostol (Cytotec) and 2 hours after insertion of misoprostol (Cytotec).

Medications

Medication Information(NOTE)*

Order with caution- less than 30 weeks GA with uterine scar, unexplained vaginal bleeding, placenta previa/vasa, fever, glaucoma, asthma, cardiac, renal, or hepatic dysfunction.(NOTE)*

Contraindications- 1. Hypersensitivity to prostaglandins.(NOTE)*

Contraindications- 2. Patient receiving oxytocin/other ripening agent.(NOTE)*

Contraindications- 3. 30 weeks or greater GA with previous uterine scar.(NOTE)*

Second Trimester of labor use order below:(NOTE)*

- ☐ **+1 Hours** miSOPROStol

- ☐ 400 mcg, Tab, VAG, q3h, Routine, (for 4 dose) (DEF)*

Comments: for fetal demise, stop administration at delivery

- ☐ 200 mcg, Tab, VAG, q4h, Routine, (for 4 dose)

Comments: for fetal demise, stop administration at delivery

Third Trimester of labor use order below:(NOTE)*

- ☐ **+1 Hours** miSOPROStol

100 mcg, Tab, VAG, prn, PRN Other, specify in Comment, Routine

Comments: to stimulate active labor

Date

Time

Physician's Signature

MD Number

*Report Legend:

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal





Physician Orders ADULT: OB Fetal Demise Plan

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

