
 Methodist Le Bonheur Healthcare		attach patient label here	
		Physician Orders ADULT	
		Order Set: Intraventricular Alteplase for Intracerebral Hemorrhage Protocol	
[R] = will be ordered			
T= Today; N = Now (date and time ordered)			
Height: _____ cm Weight: _____ kg			
Allergies:		<input type="checkbox"/> No known allergies	
<input type="checkbox"/> Medication allergy(s): _____			
<input type="checkbox"/> Latex allergy <input type="checkbox"/> Other: _____			
NOTE: The administration of Intraventricular Alteplase is restricted to Physicians ONLY and requires a nursing double check when documented on the eMAR.			
[R]	Intraventricular Alteplase for Intracerebral Hemorrhage Protocol Orders	T;N	
Patient Care			
<input type="checkbox"/>	Vital Signs	Monitor and Record T, P, R, BP Routine, q15min, For 1 hr, While EVD clamped.	
<input type="checkbox"/>	Vital Signs	Monitor and Record T, P, R, BP Routine, q 30min, For 1 hr, While EVD clamped.	
<input type="checkbox"/>	ICP Monitoring	T;N, q15min x 1 hour While EVD clamped	
<input type="checkbox"/>	ICP Monitoring	T;N, q30min x 1 hour after EVD opened	
Medications			
<input type="checkbox"/>	alteplase	1mg, injection, Device, once, STAT, Comments: Remove 5 mL of CSF from EVD prior to administration. Give 1 mg/1 mL alteplase through EVD, flush with 3 mL of sodium chloride 0.9%. Clamp EVD for 1 hour following administration. Intraventricular alteplase may be repeated in 8 hours if necessary. Dose to be drawn up in hood in IV room. DO NOT send unconstituted vial to floor.	
<input type="checkbox"/>	alteplase	2 mg, Injection, Device, once, STAT, Remove 5 mL of CSF from EVD prior to administration. Give 2 mg/2 mL alteplase through EVD, flush with 3 mL of sodium chloride 0.9%. Clamp EVD for 1 hour following administration. Dose to be drawn up in hood in IV room. DO NOT send unconstituted vial to floor.	
<input type="checkbox"/>	Sodium chloride 0.9% injectable solution	3 mL, Device, Device, once, STAT, Comments: Flush 3 mL sodium chloride 0.9% following administration of intraventricular alteplase. Dose to be sent up from Pharmacy. DO NOT use flushes available on unit. Pharmacy: send Preservative Free flush.	
Consults/Notifications			
<input type="checkbox"/>	Notify Physician-Continuous	T;N, STAT, If ICP is greater than 25 mmHg for more than 5 minutes at any point while EVD is clamped, unclamp EVD and call MD.	
Date	Time	Physician's Signature	MD Number
PT-Intraventricular Alteplase for Intracerebral Hemorrhage Protocol-23049-QM0512-Rev012219 Page1 of 1		 * 1 1 1 *	