

Ht: _____ cm

SUCROSE ADMINISTRATION FOR ANALGESIA DURING NEONATAL CIRCUMCISION

Wt · ka

TIME	PHYSICIAN'S ORDERS AND DIET	DATE & TIME	PROGRESS RECORD
			Note Progress of Case, Complications, Consultations, Change in Diagnosis, Condition on Discharge, Instructions to Patient.
	1. Prior to preparation of the operative field, nursing will		
	assess pain level using appropriate pain scale.		
	2. During preparation of the operative field, nursing will		
	administer:		
	□ Sucrose solution by repeatedly dipping pacifier in		
	24% sucrose solution 2 minutes before procedure		
	(a small pacifier dipped once holds approximately		
	0.1 ml). Monitor pain response during and after		
	procedure.		
	OR		
	□ 1 ml 24% sucrose solution by oral syringe for infants		
	less than 3000 grams 2 minutes prior to procedure.		
	Monitor pain response during and after procedure.		
	(Repeat oral syringe dose every 2 minutes as needed,		
	with a total of 3 doses per procedure.)		
	□ 2 ml 24% sucrose solution by oral syringe for infants		
	greater than or equal to 3000 grams 2 minutes prior		
	to procedure. Monitor pain response during and after		
	procedure.		
	(Repeat oral syringe dose every 2 minutes as needed,		
	with a total of 3 doses per procedure)		
	3. Document time, dose and response with each		
	administration.		
	Physician Signature:		
	Physician ID Number:		

Unapproved abbreviations – U, IU, QD, QOD, MS, MSO4, MgSO4, doses without leading zeros, doses with trailing zeros, AU, AD, AS, AL, OS, OD, OU, ug. 856-QM-OBCIRCUMSUCROSE-0106-VER3