



Physician Orders PEDIATRIC: PED GEN SURG Post Op Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- ☒ Initiate Powerplan Phase

Phase: PED GEN SURG Post Op Plan, When to Initiate: _____

PED GEN SURG Post Op Phase

Admission/Transfer/Discharge

- ☐ Return Patient to Room
T;N
- ☐ Transfer Pt within current facility

Vital Signs

- ☐ Vital Signs
Monitor and Record T,P,R,BP, q4h(std)

Activity

- ☐ Bedrest
Routine
- ☐ Bedrest w/BRP
- ☐ Out Of Bed
Up As Tolerated

Food/Nutrition

- ☐ NPO
Start at: T
- ☐ Clear Liquid Diet
Start at: T;N

Patient Care

- ☐ Advance Diet As Tolerated
- ☐ Incentive Spirometry NSG
q1h-Awake
- ☐ Nothing Per Rectum
T;N, Place sign above patient's bed-Nothing Per Rectum.
- ☐ Intake and Output
q2h(std)
- ☐ Cardiopulmonary Monitor
Monitor Type: CP Monitor, Special Instructions: Upon arrival to floor
- ☐ Discontinue CP Monitor
When ALL Criteria met: No NG-Tube, No PCA, No Chest Tube, No Sepsis Alert notified, PEWS of 0 and 24 hour post op.
- ☐ O2 Sat Spot Check-NSG
T;N, q4h(std)
- ☐ O2 Sat Continuous Monitoring NSG
q2h(std)





Physician Orders PEDIATRIC: PED GEN SURG Post Op Plan

- ☐ NGT Insert
- ☐ NGT
Suction Strength: Low Intermittent
- ☐ Indwelling Urinary Catheter Insert-Follow Removal Protocol
Reason: Postop Surgery Less Than 24 Hrs Ago, bedside gravity drainage
- ☐ Indwelling Urinary Catheter Care
Bedside gravity drainage
- ☐ Convert IV to INT/Hepwell
T;N

Respiratory Care

- ☐ Oxygen Delivery (Ped)
Special Instructions: Titrate to keep O2 sat \geq 92%, wean to room air

Continuous Infusion

- ☐ D5 1/2 NS KCl 20 mEq/L
1,000 mL, IV, Routine, mL/hr
- ☐ LR
1,000 mL, IV, Routine, Replacement fluids, Replace NGT output 1mL:1mL q4h

Medications

- ☐ **+1 Hours** acetaminophen
 - ☐ 10 mg/kg, Liq, PO, q4h, PRN Pain or Fever, Routine, For temperature greater than 38, Max Dose = 90 mg/kg/day up to 4 g/day (DEF)*
 - ☐ 10 mg/kg, Chew tab, PO, q4h, PRN Pain or Fever, Routine, For temperature greater than 38, Max Dose = 90 mg/kg/day up to 4 g/day
 - ☐ 325 mg, Tab, PO, q4h, PRN Pain or Fever, Routine, For temperature greater than 38, Max Dose = 90 mg/kg/day up to 4 g/day
- ☐ **+1 Hours** acetaminophen
10 mg/kg, Supp, PR, q4h, PRN Pain or Fever, Routine, For temperature greater than 38, Max Dose = 90 mg/kg/day up to 4 g/day
- ☐ **+1 Hours** acetaminophen-HYDROcodone 325 mg-7.5 mg/15 mL oral solution
0.2 mg/kg, Liq, PO, q4h, Pain, Routine, Max dose = 10 mg, (5 mL = 2.5 mg HYDROcodone)
- ☐ **+1 Hours** acetaminophen-HYDROcodone 325 mg-5 mg oral tablet
1 tab, Tab, PO, q4h, PRN Pain, Routine, (1 tab = 5 mg HYDROcodone)
- ☐ **+1 Hours** morphine
0.05 mg/kg, Ped Injectable, Urethral, q3h, PRN Pain, Routine, Max initial dose = 2 mg
Comments: Max initial dose = 2 mg
- ☐ **+1 Hours** ceFAZolin
25 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 3 dose), Max dose = 1 gram
- ☐ **+1 Hours** cefOXitin
30 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 3 dose), Max dose = 12 grams/day





Physician Orders PEDIATRIC: PED GEN SURG Post Op Plan

- ☐ **+1 Hours** cefTAZidime
50 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 14 day), Max dose = 2 grams/dose
- ☐ **+1 Hours** clindamycin
10 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 14 day), Max dose = 4.8 grams/day
- ☐ **+1 Hours** meropenem
20 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 14 day), Max dose = 2 grams
- ☐ **+1 Hours** metroNIDAZOLE
7.5 mg/kg, Ped Injectable, IV Piggyback, q6h, Routine, (for 14 day), Max 4grams/day
- ☐ **+1 Hours** vancomycin
10 mg/kg, Ped Injectable, IV Piggyback, q6h, Routine, (for 14 day), Max dose = 1 gram
- ☐ **+1 Hours** ondansetron
0.1 mg/kg, Ped Injectable, IV Piggyback, q8h, PRN Nausea, Routine, Max dose = 8 mg
- ☐ **+1 Hours** ketorolac
0.5 mg/kg, Ped Injectable, IV Push, q6h, Routine, (for 8 dose), Max dose = 30 mg
- ☐ **+1 Hours** famotidine
0.25 mg/kg, Injection, IV, q12h, Routine, Max Daily Dose = 40 mg/day

Laboratory

- ☐ CBC
STAT, T;N, once, Type: Blood
- ☐ CBC
Routine, T+1;0400, once, Type: Blood
- ☐ BMP
STAT, T;N, once, Type: Blood
- ☐ BMP
Routine, T+1;0400, once, Type: Blood
- ☐ Hgb & Hct
STAT, T;N, once, Type: Blood
- ☐ Hgb & Hct
Routine, T+1;0400, once, Type: Blood

Diagnostic Tests

- ☐ Chest 1 VW
T;N, Stat, Portable

Consults/Notifications/Referrals

- ☐ Notify Physician-Continuing
Notify: Resident on call, Notify For: For temperature greater than 38.0 degrees, persistent vomiting, urinary output less than 1mL/kg/hr over four hours, or signs/symptoms of surgical site infection (erythema/blood/pus)
- ☐ Physician Group Consult
- ☐ Physician Consult





Physician Orders PEDIATRIC: PED GEN SURG Post Op Plan

☐ PICC Line Insert
Reason for Consult: PICC Line Insertion

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

***Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

