



Physician Orders ADULT: General Surgery Postop Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- ☒ Initiate Powerplan Phase

Phase: General Surgery Postop Phase, When to Initiate: _____

General Surgery Postop Phase

Admission/Transfer/Discharge

- ☐ Return Patient to Room
T;N
- ☐ Transfer Pt within current facility

Vital Signs

- ☐ Vital Signs
P, R, BP q30 min x 2; P, R, BP, q1hr x 2; P, R, BP q 4hr x 6, Temperature on admission to unit and q4hr x 6
- ☐ Vital Signs
Monitor and record T, P, R, BP q2hr x 8, then q shift

Activity

- ☐ Dangle At Bedside
twice per shift
- ☒ Up To Chair
T+1;N
- ☐ Ambulate
bid

Food/Nutrition

- ☐ NPO
- ☐ Clear Liquid Diet
Start at: T;N

Patient Care

- ☒ Incentive Spirometry NSG
q2h-Awake
- ☐ Intake and Output
Routine, q4h(std), Record Intake & Output
- ☐ Cough and Deep Breathe
- ☐ Measure Circumference
T;N
- ☐ Abdominal Binder Apply
- ☒ In and Out Cath
PRN if unable to void x 1





Physician Orders ADULT: General Surgery Postop Plan

- ☐ Irrigate
Nasogastric Tube, Routine, q4h(std)
- ☐ NGT
Suction Strength: Low Intermittent
- ☐ NGT
Do not reposition NG tube
- ☐ NGT
Reposition NG if not draining
- ☐ Drain Care
q8h(std), Empty and record JP output
- ☐ Dressing Care
T;N, Action: Change, PRN, Change drain dressing PRN

Respiratory Care

- ☐ O2-Nasal Cannula
Routine, 2 L/min, Special Instructions: titrate to keep O2 sat \geq 92%

Continuous Infusion

- ☐ D5W with NaCl 0.45%
1,000 mL, IV, Routine, 45 mL/hr
- ☐ Lactated Ringers Injection
1,000 mL, IV, Routine, 75 mL/hr

Medications

- ☐ VTE Other SURGICAL Prophylaxis Plan(SUB)*
- ☐ **+1 Hours** famotidine
20 mg, Injection, IV Push, q12h, Routine
Pick ONE of the PCA plans:(NOTE)*
- ☐ PCA - MorPHINE Protocol Plan (Adult)(SUB)*
- ☐ PCA - HYDROmorphine Protocol Plan (Adult)(SUB)*
Pick ONE of the following for Breakthrough pain:(NOTE)*
- ☐ **+1 Hours** morphine
1 mg, Injection, IV Push, q4h, PRN Pain, Breakthrough, Routine
- ☐ **+1 Hours** HYDROmorphine
0.5 mg, Injection, IV Push, q4h, PRN Pain, Breakthrough, Routine
- ☐ **+1 Hours** ondansetron
4 mg, Injection, IV Push, q4h, PRN Nausea, Routine
Comments: Give first to treat nausea
- ☐ **+1 Hours** prochlorperazine
5 mg, Injection, IV Push, q4h, PRN Nausea
Comments: Give if no relief from ondansetron





Physician Orders ADULT: General Surgery Postop Plan

- ☐ **+1 Hours** ceFAZolin
2 g, IV Piggyback, IV Piggyback, q8h, Routine, (for 2 dose)
- ☐ **+1 Hours** vancomycin
15 mg/kg, Injection, IV Piggyback, q12h, Routine, (for 1 dose)
- ☐ **+1 Hours** metroNIDAZOLE
500 mg, IV Piggyback, IV Piggyback, q8h, Routine, (for 2 dose)
- ☐ **+1 Hours** piperacillin-tazobactam
4.5 g, IV Piggyback, IV Piggyback, q6h, Routine, (for 3 dose)
If Allergic To Penicillin:(NOTE)*
- ☐ **+1 Hours** clindamycin
600 mg, IV Piggyback, IV Piggyback, q8h, Routine, (for 3 dose)
- ☐ **+1 Hours** gentamicin
1.5 mg/kg, IV Piggyback, IV Piggyback, q8h, Routine, (for 3 dose)

Laboratory

- ☐ CBC w/o Diff
Routine, T+1;0400, once, Type: Blood
- ☐ Hct
Routine, T+1;0400, once, Type: Blood
- ☐ CMP
Routine, T+1;0400, once, Type: Blood
- ☐ BMP
Routine, T+1;0400, once, Type: Blood
- ☐ PT/INR
Routine, T+1;0400, once, Type: Blood
- ☐ PTT
Routine, T+1;0400, once, Type: Blood
- ☐ Type and Crossmatch PRBC
Routine, T+1;0400, Type: Blood
- ☐ Hold PRBC
Routine, T+1;0400, Reason: On Hold for OR
- ☐ Urinalysis
Routine, T+1;0400, once, Type: Urine, Nurse Collect
- ☐ Urinalysis w/Reflex Microscopic Exam
Routine, T+1;0400, once, Type: Urine, Nurse Collect

Diagnostic Tests

- ☐ Chest 1VW Frontal
T;N, Portable

Consults/Notifications/Referrals





Physician Orders ADULT: General Surgery Postop Plan

- ☐ Notify Physician-Once
Notify For: Notify of room number upon arrival to unit
- ☐ Notify Physician-Continuing
Notify For: if patient unable to void
- ☐ Notify Physician For Vital Signs Of
BP Systolic > 170, BP Diastolic > 110, BP Systolic < 95, BP Diastolic < 55, Heart Rate > 110, Heart Rate < 60, Resp Rate > 24, Resp Rate < 10, Urine Output < 250 mL/8hrs

Date	Time	Physician's Signature	MD Number

***Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

