



## LEB ED Children's Intestinal Rehab Center LeBonheur (CIRCLE) Fever Plan

**LEB ED CIRCLE Fever Phase****Non Categorized**

Criteria: Patient goes to CIRCLE clinic, has a central line and fever greater than 100.4. (NOTE)\*

R Powerplan Open

Notify Physician-Once

*Notify: On Call GI Physician, Notify For: Upon arrival to ED*

**Vital Signs**

Vital Signs

*Monitor and Record T,P,R,BP, q30min*

**Patient Care**

Central Line May Use

*Routine, May use Central Line for: No Limits*

Bedside Glucose Nsg

Cardiopulmonary Monitor

*T;N, Monitor Type: CP Monitor, Special Instructions: q12h*

**Nursing Communication**

Nursing Communication

*Discontinue home TPN on arrival.*

Nursing Communication

*Consult MD for D10 1/2NS fluid rate.*

**Respiratory Care**

O2 Sat-Continuous Monitoring (RT)

*now*

**Continuous Infusion**

D10 NS (Pediatric) (IVS)\*

Dextrose 10% in Water

1,000 mL, IV, STAT

*Comments: Consult MD for Fluid rate.*

sodium chloride

154 mEq

**Medications**

acetaminophen

*15 mg/kg, Liq, PO, q6h, PRN Fever, STAT, Max single dose = 650 mg*

cefTAZidime

*50 mg/kg, Ped Injectable, Central IV, once, STAT, Max single dose = 2,000 mg*

*Comments: To Be Infused First*

vancomycin

*15 mg/kg, Injection, Central IV, once, STAT, ( infuse over 1 hr )*

**Laboratory**

CBC with Diff

*STAT, T;N, once, Type: Blood*

CMP

*STAT, T;N, once, Type: Blood*

Blood Culture Pediatric Plan(SUB)\*

**LEB ED CIRCLE Fever PHASE 2 - MD ONLY****Continuous Infusion**

Sodium Chloride 0.9% Bolus

*20 mL/kg, Injection, IV, once, STAT, Max dose =1 liter*

**Medications**

**+1 Hours** linezolid

*10 mg/kg, Ped Injectable, Central IV, once, STAT, Max single dose = 600 mg*

Place order below instead of ceftazidime if shock or has had Enterobacter, Serratia, Citrobacter, Morganella or another ceftazidime resistant line sepsis in the previous year:(NOTE)\*





**LEB ED Children's Intestinal Rehab Center LeBonheur (CIRCLE) Fever Plan**

- meropenem  
*20 mg/kg, Ped Injectable, Central IV, once, STAT, Max single dose = 1,000 mg*  
*Comments: To Be Infused First*
- Place order below if history of Candida line sepsis or shock:(NOTE)\*
- +1 Hours** micafungin  
*2 mg/kg, Ped Injectable, Central IV, once, STAT, ( infuse over 1 hr ), Max single dose = 100mg*

**Laboratory**

Please order below for abdominal symptoms :(NOTE)\*

- Amylase Level  
*STAT, T;N, once, Type: Blood*
- Amylase Level LeBonheur Germantown  
*STAT, T;N, once, Type: Blood*
- Lipase Level  
*STAT, T;N, once, Type: Blood*
- Lipase Level LeBonheur Germantown  
*STAT, T;N, once, Type: Blood*

Place orders below for symptoms of shock :(NOTE)\*

- Cortisol Level  
*STAT, T;N, once, Type: Blood*
- PT/INR  
*STAT, T;N, once, Type: Blood*
- PTT  
*STAT, T;N, once, Type: Blood*

Strong consideration should be given to ordering a Cath UA and Culture on all infants less than 12 months old with no readily identifiable cause for their fever.(NOTE)\*

- Urine Culture  
*STAT, T;N, Specimen Source: Urine, Clean Catch, Nurse Collect*
- Urinalysis w/Reflex Microscopic Exam  
*STAT, T;N, once, Type: Urine, Nurse Collect*

**Diagnostic Tests**

- Chest 1 VW  
*T;N, Reason For Exam Respiratory Symptoms, Stat, Portable*
- Chest 2 Views  
*T;N, Reason For Exam Respiratory Symptoms, Stat, Portable*

**Date**

**Time**

**Physician's Signature**

**MD Number**

**\*Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

