

attach patient label here



Physician Orders - Adult
Heparin Induced Thrombocytopenia (HIT) Protocol Orders

[X or R] = will be ordered unless marked out.

T= Today; N = Now (date and time ordered)

Height: _____ cm Weight: _____ kg

Allergies: ☐ No known allergies

☐ Medication allergy(s): _____

☐ Latex allergy ☐ Other: _____

Medications

NOTE: Select either Bivalirudin, Argatroban, or Fondaparinux Orders below:

NOTE: Argatroban preferred when CrCl is less than 15 mL/min or on dialysis.

NOTE: Fondaparinux is contraindicated when CrCl is less than 30 mL/min

☐ Bivalirudin Orders

☐ Argatroban Orders

☐ Fondaparinux Orders

CareSets/Protocols/PowerPlans

[R] Heparin Induced Thrombocytopenia (HIT) T;N



Physician Orders: Adult
Heparin Induced Thrombocytopenia (HIT) Protocol: Bivalirudin
Orders

[X or R] = will be ordered unless marked out.

T= Today; N = Now (date and time ordered)

Height: _____ cm Weight: _____ kg

Allergies:		<input type="checkbox"/> No known allergies
<input checked="" type="checkbox"/> Medication allergy(s): Heparin Low Molecular Weight Heparins _____		
<input type="checkbox"/> Latex allergy		<input type="checkbox"/> Other: _____
Patient Care		
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Bivalirudin: Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Adjust rate of bivalirudin based upon Bivalirudin Dose Adjustment Instructions (Ref Text)(pg. 3 for paper)
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Document the initiation, the rate, rate changes, and discontinuation on the HIT Protocol Flow Record.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Document the time of aPTT lab draw and result on the HIT Protocol Flow Record.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Discontinue daily CBC and aPTT when bivalirudin is discontinued.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Bivalirudin-Validate that heparin is documented as an allergy
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Discontinue orders for any heparin or LMWH, including heparin flushes or locks.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: If heparin or LMWH is ordered, contact prescriber to inform of Heparin allergy/HIT so that alternative anticoagulation can be ordered.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Label all IV sites/catheters "NO HEPARIN".
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: No Heparin coated needles for ABGs.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Place order for Partial Thromboplastin Time (PTT) to be collected 2 hours after the start of Bivalirudin
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: Place order for and Draw PTT 2 hours after any rate change on Bivalirudin infusion
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Bivalirudin: If any two sequential aPTT results exceed 75 seconds while on bivalirudin protocol, notify Pharmacist.





Physician Orders: Adult
Heparin Induced Thrombocytopenia (HIT) Protocol: Bivalirudin
Orders

[X or R] = will be ordered unless marked out.

T= Today; N = Now (date and time ordered)

Medications	
Bivalirudin Maintenance	
[]	.Bivalirudin (HIT) Protocol Reference Text See Ref. Text, N/A, N/A, Routine
NOTE: If CrCl less than 10 mL/min or conv HD, place Bivalirudin order below:	
[]	bivalirudin (additive) 250 mg + Sodium Chloride 0.9% 250 mL 250 mL, IV, Routine,T;N, Titrate, Comment: Start rate at 0.02 mg/kg/hr, titrate per protocol. BIVALIRUDIN DOSE ADJUSTMENT INSTRUCTIONS (Use Standard Concentration 1 mg / mL) aPTT (seconds) Dose Adjustment /Monitoring
NOTE: If CrCl between 10-29 mL/min or CRRT, place Bivalirudin order below:	
[]	bivalirudin (additive) 250 mg + Sodium Chloride 0.9% 250 mL 250 mL, IV, Routine,T;N, Titrate, Comment: Start rate at 0.05mg/kg/hr, titrate per protocol. BIVALIRUDIN DOSE ADJUSTMENT INSTRUCTIONS (Use Standard Concentration 1 mg / mL) aPTT (seconds) Dose Adjustment /Monitoring
NOTE: If CrCl between 30-59 mL/min, place Bivalirudin order below:	
[]	bivalirudin (additive) 250 mg + Sodium Chloride 0.9% 250 mL 250 mL, IV, Routine,T;N, Titrate, Comment: Start rate at 0.08 mg/kg/hr, titrate per protocol BIVALIRUDIN DOSE ADJUSTMENT INSTRUCTIONS (Use Standard Concentration 1 mg / mL) aPTT (seconds) Dose Adjustment /Monitoring
NOTE: If CrCl greater than 60mL/min, place Bivalirudin order below:	
[]	bivalirudin (additive) 250 mg + Sodium Chloride 0.9% 250 mL 250 mL, IV, Routine,T;N, Titrate, Comment: Start rate at 0.15 mg/kg/hr, titrate per protocol BIVALIRUDIN DOSE ADJUSTMENT INSTRUCTIONS (Use Standard Concentration 1 mg / mL) aPTT (seconds) Dose Adjustment /Monitoring
Laboratory	
[]	Partial Thromboplastin Time (PTT) T;N, STAT, once, Type: Blood, Order Comment: draw prior to start of Bivalirudin infusion
[]	CBC Routine, T+1;0400, qam, Type: Blood
Diagnostic Tests	
NOTE: If not already performed, order bilateral ultrasound venous doppler below:	
[]	US Ext Lower Ven Doppler W Compress Bil T;N, Reason for Exam: DVT (Deep Vein Thrombosis), Routine, Stretcher, Comment: Heparin Allergy
Consults/Notifications	
[]	Pharmacy Consult-HIT Bivalirudin dosing T;N, Notify Clinical Pharmacy Specialist for initiation and daily follow-up of bivalirudin
[]	Pharmacy Consult-Warfarin Dosing T;N, qam
CareSets/Protocols/PowerPlans	
[R]	Bivalirudin (HIT) Protocol Orders.

Date

Time

Physician's Signature

MD Number





Physician Orders-Adult
Heparin Induced Thrombocytopenia (HIT) Protocol:
Argatroban Orders

attach patient label here

[X or R] = will be ordered unless marked out.

T= Today; N = Now (date and time ordered)

Height: cm Weight: kg

Allergies: ☐ No known allergies

☐ Medication allergy(s):

☐ Latex allergy ☐ Other:

Nursing Communication

[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: Validate that heparin is documented as an allergy
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: Discontinue orders for any heparin or LMWH, including heparin flushes or locks
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: If heparin or LMWH is ordered, contact prescriber to inform of Heparin allergy/HIT so that alternative anticoagulation can be ordered
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: Label all IV sites/catheters "NO HEPARIN"
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: No heparin coated needles for ABG's
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: Place order for Partial Thromboplastin Time (PTT) to be collected 2 hours after the start of argatroban continuous infusion.
[X]	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Argatroban: Draw PTT 2 hours after any rate change on Argatroban infusion
[X]	Nursing Communication	sequential PTT results exceed 90seconds while on Argatroban protocol, notify Pharmacist

Argatroban Maintenance Medications

NOTE: Argatroban preferred when CrCl is less than 15 mL/min or on dialysis.

NOTE: If total bilirubin is equal to or less than 1.5mg/dL, place argatroban order below:

[]	argatroban (Argatroban 50 mg/50mL Injectable Solution)	50 mg / 50 mL, IV, per protocol, Titrate, Comment: initial infusion is 1 mcg/kg/min.
		Titrate per protocol below:
		Comment: PTT(sec) Rate Adjustment Draw PTT

NOTE: If total bilirubin is greater than 1.5mg/dL, place argatroban order below:

[]	argatroban (Argatroban 50 mg/50mL Injectable Solution)	50 mg / 50 mL, IV, per protocol, Titrate, Comment: initial infusion is 0.5.mcg/kg/min
		Titrate per protocol below:
		PTT(sec) Rate Adjustment Draw PTT
		>90 Stop infusion. 2 hours post rate change
[R]	Sodium Chloride 0.9%	1,000 mL, IV, Routine, 20 mL/hr (infuse over 50 hr), Comment: To be run to keep line open when argatroban rate falls below 10 mL/hr.

Laboratory

[]	CBC	Routine, T+1;0400, qam, Type: Blood
[]	Partial Thromboplastin Time (PTT)	T;N, STAT, once, Type: Blood
[]	Partial Thromboplastin Time (PTT)	Routine, T+1;0400, qam, Type: Blood

Diagnostic Tests

[]	US Ext Lower Ven Doppler W Compress Bil	T;N, Reason for Exam: DVT (Deep Vein Thrombosis), Routine, Stretcher, Comment: Heparin Allergy
-----	---	--

Consults/Notifications

[]	Pharmacy Consult-Warfarin Dosing	Routine, qam
[]	Pharmacy Consult-DTI Dosing-argatroban	Routine, qam, DTI - Argatroban dosing

CareSets/Protocols/PowerPlans

[R]	Argatroban (HIT) Protocol Orders.
-----	-----------------------------------

Date

Time

Physician's Signature

MD Number



* 1 1 1 *



Physician Orders - Adult
Heparin Induced Thrombocytopenia (HIT) Protocol:
Fondaparinux Orders

attach patient label here

[X or R] = will be ordered unless marked out.

T= Today; N = Now (date and time ordered)

Height: _____ cm Weight: _____ kg

Allergies: ☐ No known allergies

☐ Medication allergy(s): _____

☐ Latex allergy ☐ Other: _____

Nursing Communication

<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Validate that heparin is documented as an allergy
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Discontinue orders for any heparin or LMWH, including heparin flushes or locks
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: If heparin or LMWH is ordered, contact prescriber to inform of Heparin allergy/HIT so that alternative anticoagulation can be ordered
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Fondaparinux Label all IV sites/catheters "NO HEPARIN"
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Fondaparinux No heparin coated needles for ABG's
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Discontinue any active orders for Bivalirudin, Argatroban or Heparin. If patient was previously on Bivalirudin/Argatroban, obtain PTT every 4 hours. Do not start Fondaparinux until PTT less than 45 seconds.
<input checked="" type="checkbox"/>	Nursing Communication	Heparin Induced Thrombocytopenia (HIT) Protocol Fondaparinux: Page Clinical Pharmacist for Fondaparinux initiation and daily follow-up.

Fondaparinux for NON Acute HIT and weight greater than 50kg

<input type="checkbox"/>	fondaparinux	2.5 mg, Injection, Subcutaneous, QDay, Routine, T;N, Comment: Do not start until PTT is less than 45 seconds if patient was previously on Bivalirudin, Argatroban or heparin.
--------------------------	--------------	---

Fondaparinux for Acute HIT or thrombosis present

NOTE: If weight less than 50kg place order below:

<input type="checkbox"/>	fondaparinux	5 mg, Injection, Subcutaneous, QDay, Routine, T;N, Comment: Do not start until PTT is less than 45 seconds if patient was previously on Bivalirudin, Argatroban or Heparin.
--------------------------	--------------	---

NOTE: If weight is 50-100kg place order below:

<input type="checkbox"/>	fondaparinux	7.5 mg, Injection, Subcutaneous, QDay, Routine, T;N, Comment: Do not start until PTT is less than 45 seconds if patient was previously on Bivalirudin, Argatroban or Heparin.
--------------------------	--------------	---

NOTE: If weight is greater than 100kg place order below:

<input type="checkbox"/>	fondaparinux	10 mg, Injection, Subcutaneous, QDay, Routine, T;N, Comment: Do not start until PTT is less than 45 seconds if patient was previously on Bivalirudin, Argatroban or Heparin.
--------------------------	--------------	--

Laboratory

<input type="checkbox"/>	CBC w/o Diff	Routine, T+1;0400, qam, Type: Blood
--------------------------	--------------	-------------------------------------





Physician Orders - Adult

Title: Direct Thrombin Inhibitor (DTI) Protocol Fondaparinux Orders

attach patient label here

[X or R] = will be ordered unless marked out.
T= Today; N = Now (date and time ordered)

Diagnostic Tests

NOTE: If not already performed, order bilateral ultrasound venous doppler below:

[]	US Ext Lower Ven Doppler W Compress Bil	T;N, Reason for Exam: DVT (Deep Vein Thrombosis), Routine, Stretcher, Comment: Heparin Allergy
-----	--	---

Consults/Notifications

[X]	Consult Clinical Pharmacist	Start at: T;N, Special Instructions: Discontinue Bivalirudin, Argatroban or Heparin if patient is currently receiving.
[X]	Pharmacy Consult-DTI Fondaparinux dosing	Routine, once, DTI - Fondaparinux dosing
[X]	Pharmacy Consult-Warfarin Dosing	Routine qam

Date

Time

Physician's Signature

MD Number



PHYSICIAN'S ORDERS

HT: _____ cm

WT: _____ kg

DATE: _____

Allergies: Heparin Low-Molecular Weight Heparins

TIME: _____

HEPARIN-INDUCED THROMBOCYTOPENIA (DTI) PROTOCOL –BIVALIRUDIN**Orders completed by Nursing**

1. Page Clinical Pharmacy Specialist / Coordinator for initiation and daily follow-up.
2. Order CBC without differential DAILY.
3. Draw baseline aPTT prior to infusion.
4. STAT aPTT 2 hours after the start of the continuous infusion and 2 hours after any rate change.
5. Stop all heparin or low-molecular weight heparin, including flushes or locks.
6. Label all IV sites or catheters as "NO HEPARIN"
7. Adjust rate of infusion based upon *Bivalirudin Dose Adjustment Instructions*.

BIVALIRUDIN DOSE ADJUSTMENT INSTRUCTIONS (Use Standard Concentration 1 mg / mL)	
aPTT (seconds)	Dose Adjustment /Monitoring
Greater than 75	Stop infusion for 1 hour and then restart at 50% slower rate. (new rate=current rate/2) (Reminder - Draw aPTT 2 (two) hours after each rate change)
45-75	Continue at current rate. Draw aPTT in AM
Less than 45	Increase infusion rate by 20% . (new rate=current rate x 1.2) (Reminder - Draw aPTT 2 (two) hours after each rate change)

8. Document the initiation, the rate, rate changes, and discontinuation on the *HIT Protocol Flow Record*
9. Document the time of aPTT lab draw and result on the *HIT Protocol Flow Record*
10. Discontinue daily CBC and aPTT when bivalirudin is discontinued
11. If any two sequential aPTTs exceed 75 seconds, page the Clinical Pharmacy Specialist On-Call/Coordinator at _____.

Orders for Pharmacist

1. Order bilateral lower extremity ultrasound for DVT if not already done
2. Discontinue active orders for any heparin or LMWH and add to allergy list
3. Calculate CrCl using Cockcroft-Gault equation

Initial Maintenance Infusion (250mg / 250ml NS or D5W)	
CrCl (ml/min)	Dose (based on actual body weight)
> 60	0.15 mg/kg/hr
30-59	0.08 mg/kg/hr
10-29 or CRRT	0.05 mg/kg/hr
< 10 or conv HD	0.02 mg/kg/hr

4. Enter initial infusion rate _____ mL/hr

Orders for Physician

- ☐ Warfarin Dosing Service to follow & begin anticoagulation with warfarin after platelet count recovery & when physician specifies.
- ☐ Do not consult Warfarin Dosing Service. MD to manage warfarin.

Warfarin Management Recommendations (NOT ORDERS)

1. Do not start warfarin until platelets > 150,000 / mm³
2. Use doses no greater than 5 mg to initiate warfarin therapy
3. Minimum of 5 days of overlap with bivalirudin and warfarin
4. **NOTE:** Bivalirudin slightly elevates the INR *in vitro*; therefore, overlap with warfarin until INR greater than 3
5. Once INR greater than 3 for 2 consecutive days, stop bivalirudin

Physician Signature:
SignaturePhysician number:
Date

Date/Time: