



## Physician Orders - ADULT

attach patient label here

### VTE- Cesarean Section Surgical Prophylaxis Plan

[R] = will be ordered

T= Today; N = Now (date and time ordered)

Height: \_\_\_\_\_ cm Weight: \_\_\_\_\_ kg

Allergies: \_\_\_\_\_ ☐ No known allergies

☐ Medication allergy(s): \_\_\_\_\_

☐ Latex allergy ☐ Other: \_\_\_\_\_

**NOTE: Bleeding Risk Factor Assessment criteria is listed below VTE orders.**

#### VTE Orders

**NO Additional Medical VTE Risk Factors Present and NO Contraindications to SCDs:**

☐ Sequential Compression Device T;N, Apply To: Lower Extremities, Comment: Bleeding Risks Present Apply

**NO Additional Medical VTE Risk Factors Present and Contraindications to SCDs:**

☐ Ambulate T;N

**NO Bleeding Risks Present AND One or More Medical VTE Risk Factors Present:**

☐ heparin 5,000 units, Injection, Subcutaneous, q8h, Routine, T;N +720, Do not adjust time of first dose as scheduled by pharmacy.

**OR**

☐ enoxaparin 40 mg, Injection, Subcutaneous, Qday, Routine, T;N + 720, Comment: first dose should be timed to be given at least 12 hours post op. If CrCl less than 30 mL/min, pharmacy to adjust dose to 30mg SQ Qday. Do not adjust time of first dose as scheduled by pharmacy.

**AND BOTH CBCs:**

☐ CBC w/o Diff Routine, T;N, once, Type: Blood

☐ CBC w/o Diff Time Study, T+2:0400, Every Other Day, Type: Blood

#### Do Not Administer VTE Prophylaxis

☐ Contraindication-VTE Prophylaxis T;N, Reason: Patient has bleeding risk for anticoagulants, and SCDs are contraindicated. Consider early ambulation.

**BLEEDING RISK FACTOR ASSESSMENT:** This is a partial list of bleeding risk factors. Clinicians are advised to consider other risk factors or conditions that may predispose patients to DVT/PE. Check all that may apply:

☐ Patient already receiving anticoagulation therapy with warfarin, heparin, fondaparinux, enoxaparin or other anticoagulation therapy

☐ Active bleeding

☐ INR greater than 1.5 and patient **NOT** on warfarin therapy

☐ INR greater than 2 and patient **ON** warfarin therapy

☐ Solid Organ and Bone Marrow Transplant patients with platelet count less than 100,000

☐ Platelet count less than 50,000 (applies to patients with no history of transplant procedures)

☐ Solid organ transplant during this episode of care **OR** within 30 days of admission

☐ Documented bleeding or Coagulopathy disorder

☐ Hemorrhagic Stroke within 6 weeks of admission

☐ Severe Uncontrolled Hypertension

☐ Recent Intraocular or Intracranial surgery

☐ Vascular Access or Biopsy sites inaccessible to hemostatic control

☐ Recent Spinal Surgery

☐ Epidural or Spinal Catheter

☐ Pregnancy, Possible Pregnancy or Postpartum (to include up to 6 weeks post partum)

☐ Heparin Induced Thrombocytopenia (HIT)

☐ Heparin allergy or pork allergy

☐ No Bleeding Risk Factors exists

Date \_\_\_\_\_

Time \_\_\_\_\_

Physician's Signature \_\_\_\_\_

MD Number \_\_\_\_\_

VTE C-Sec Surgical Prophylaxis-25013 -  
QM-0510 Ver 2-121510

