



### Physician Orders PEDIATRIC: LEB ED Intrapartum/Postpartum Plan

#### Triage Standing Intrapartum/Postpartum

##### Non Categorized

Criteria: Patients less than 18 years of age with acute ongoing bleeding in the intra or post partum period.(NOTE)\*

##### Vital Signs

- Vital Signs  
*T;N, Stat, BP, HR, RR q15min, Temp q1hr*

##### Food/Nutrition

- NPO  
*Start at: T;N, Instructions: NPO except for ice chips | NPO except for medications, and popsicles*

##### Patient Care

- IV Insert/Site Care LEB  
*T;N, Stat, Preferred Gauge: 18G, or 20 gauge catheter*

##### Continuous Infusion

- D5LR  
*1,000 mL, IV, STAT, 125 mL/hr ( infuse over 8 hr )*

##### Laboratory

- CBC  
*STAT, T;N, once, Type: Blood*
- PT/INR  
*STAT, T;N, once, Type: Blood*
- PTT  
*STAT, T;N, once, Type: Blood*
- Fibrinogen Level  
*STAT, T;N, once, Type: Blood*
- Type and Screen Pediatric  
*STAT, T;N, Type: Blood*
- BMP  
*STAT, T;N, once, Type: Blood*
- Ionized Calcium Pediatric  
*STAT, T;N, once, Type: Blood*

##### Consults/Notifications/Referrals

- Notify Physician-Continuing  
*T;N, Notify: ED Physician, Pulse greater than 110 beats/minute; systolic BP less than 90 or greater than 160 mm/Hg; diastolic BP greater than 110 mm/Hg; temperature greater than 39 degrees Celsius; heavy vaginal bleeding (soaking two or more pads per hour)*

#### LEB ED Intrapartum/Postpartum Phase 2

##### Admission/Transfer/Discharge

- Transfer Pt within current facility





**Physician Orders PEDIATRIC: LEB ED Intrapartum/Postpartum Plan**

T;N

**Patient Care**

- In and Out Cath  
*T;N, Stat, For 2 times, PRN, for inability to void*
- Foley Insert-Follow Removal Protocol  
*T;N, If still unable to void after 2 catheterizations*
- Cold Apply  
*T;N, Perineum, for swelling/discomfort*

**Continuous Infusion**

NOTE: After delivery of placenta(NOTE)\*

- oxytocin 30 units in D5NS 500 mL (IVS)\*  
Dextrose 5% with 0.9% NaCl  
*500 mL, IV, STAT, (for 1 dose), 150 mL/hr, To be infused over 3.3 hours*  
oxytocin(additive)  
*30 units*

**Medications**

- Al hydroxide/Mg hydroxide/simethicone  
*30 mL, Oral Susp, PO, q3h, PRN Heartburn, For Indigestion or Gas*
- ibuprofen  
*800 mg, Tab, PO, q8h, PRN Other, specify in Comment, Routine, Cramping*
- acetaminophen
  - 650 mg, Tab, PO, q6h, PRN Pain or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day (DEF)\**
  - 325 mg, Tab, PO, q4h, PRN Pain or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day*
- acetaminophen-oxyCODONE 325 mg-5 mg oral tablet  
*2 tab, Tab, PO, q4h, PRN Pain, Moderate (4-7), Routine, (for 3 day)*
- HYDROMorphone  
*1 mg, Injection, IV Push, q4h, PRN Pain, Severe (8-10), Routine, (for 3 day)*
- ondansetron  
*8 mg, Ped Injectable, IV Push, q8h, PRN Nausea/Vomiting, Routine*
- NOTE: After delivery of placenta(NOTE)\*
- oxytocin  
*20 units, Injection, IM, once, STAT, Us if NO IV Access*

**Consults/Notifications/Referrals**

- Notify Physician-Once  
*T;N, of room number on arrival to unit*
- Physician Group Consult  
*T;N, Stat*

**LEB ED Postpartum Bleeding Plan Phase 3**





Physician Orders PEDIATRIC: LEB ED Intrapartum/Postpartum Plan

Patient Care

NOTE: To be initiated only after routine intrapartum oxytocin infusion is complete and when an emergent or urgent hemorrhagic situation occurs that requires multiple units and/or components of blood (cross-matched if ready or red tagged if blood sample has not been matched). This must be ordered by the ED staff physician.(NOTE)\*

Medications

NOTE: If bleeding continues post-oxytocin infusion, administer:(NOTE)\*

- methylergonovine
0.2 mg, Injection, IM, q2h, STAT, (for 2 dose )
Comments: May repeat q2hr up to 2 total doses, Administer IM only, rotating injection site with each dose.

NOTE: If bleeding persists post-METHergine administration and patient exhibits signs/symptoms of refractory postpartum uterine bleeding, administer:(NOTE)\*

- carboprost
250 mcg, Injection, IM, q15min, STAT, (for 2 dose )
Comments: May repeat q15min up to 2 total doses. Administer IM only, rotating injection site with each dose

Date Time Physician's Signature MD Number

\*Report Legend:

- DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a sub phase, see separate sheet
R-Required order

