



attach patient label here

**Physician Orders ADULT****Order Set: PICC Line Insertion Protocol Orders (PICC Nurse ONLY)**

[R] = will be ordered

T= Today; N = Now (date and time ordered)

Height: \_\_\_\_\_ cm Weight: \_\_\_\_\_ kg

<b>Allergies:</b>		<input type="checkbox"/> No known allergies
<input type="checkbox"/> Medication allergy(s): _____		
<input type="checkbox"/> Latex allergy <input type="checkbox"/> Other: _____		
Criteria for use: request for PICC nurse to place a PICC line.		
<b>Patient Care</b>		
<input type="checkbox"/>	Central Line	
<input type="checkbox"/>	PICC Line Insertion Protocol Orders	
<input type="checkbox"/>	PICC Line Care	T;N, Wednesday, Change dressing every Wednesday and PRN if dressing becomes loose, soiled, or moist.
<input type="checkbox"/>	Heat Apply	T;N, PRN Apply to Line Site, Place heat to PICC Line insetion site 3-4 times daily if site becomes red, tender, or swollen.
<input type="checkbox"/>	Measure Circumference	T;N, Qday, Of: Arm, Measure PICC arm circumference midway between elbow and axilla Qday for first 3 days after insertion.
<input type="checkbox"/>	No BP or Venipunctures	T;N in PICC Line arm. Place sign above patient's bed
<input type="checkbox"/>	Instruct/Educate	T;N, Instruct patient, Topic:PICC Catheter/Line risks and benefits
<b>Nursing Communication</b>		
<input type="checkbox"/>	Nursing Communication	T;N, Label front of chart "PICC in ___ arm to alert other departments."
<input type="checkbox"/>	Nursing Communication	T;N PICC Nurse must confirm placement of PICC prior to use.
<input type="checkbox"/>	Nursing Communication	T;N, change all tubing prior to using the PICC Line for the first time.
<b>Medications</b>		
<input type="checkbox"/>	Sodium Chloride 0.9% Flush	10 ml, Injection, IV Push, q8h, Routine, T;N, (PICC/CVL per lumen)
<input type="checkbox"/>	Sodium Chloride 0.9% Flush	20 ml, Injection, IV Push, prn, PRN, Other, specify in Comment,Comment: Use for blood administration and line draws. Routine, T;N, (PICC/CVL per lumen)
<input type="checkbox"/>	Lidocaine (lidocaine 1% inj)	3ml, Injection, Infiltration, once Routine, T;N
<b>Diagnostic</b>		
<input type="checkbox"/>	Chest 1VW Frontal	T;N, Reason for Exam: Line Placement, STAT, Portable, Comment: to be done upon completion of PICC Line Insertion to confirm PICC Line Placement
<b>Consults/Notifications</b>		
<input type="checkbox"/>	Notify Physician-Once	T;N, Call nephrologist prior to PICC line placement if end stage renal disease or if there is a nephrologist consulted on the case
<input type="checkbox"/>	Notify Physician-Continuing	T;N, Notify: PICC Line ordering MD, for PICC site redness, tenderness, edema above site, excessive bleeding at exit site, chest, neck, or ear pain, numbness or tingling of affected arm or hand.

