



attach patient label here

Physician Orders ADULT

Order Set: FCR

Diagnosis : CLL

Height: _____ cm	Weight: _____ kg	Cycle: _____ Of : _____		
Actual BSA: _____ m ²	Treatment BSA: _____ m ²	Day/Wk: _____ Freq: _____		
Allergies:				
<input type="checkbox"/> No known allergies				
<input type="checkbox"/> Medication allergy(s): _____				
<input type="checkbox"/> Latex allergy <input type="checkbox"/> Other: _____				
Patient Care				
<input type="checkbox"/> Nursing Communication	T;N, Do not exceed a treatment BSA of _____ m ²			
<input type="checkbox"/> Nursing Communication	T;N, May hold hydration during chemotherapy infusion			
Continuous Infusions				
Pre Hydration				
<input type="checkbox"/> Normal Saline	1,000 mL, IV, Routine, _____ mL/hr , Start 4 hours prior to chemotherapy and continue for at least 24 hours after chemotherapy is complete			
Medications				
Pre Medication				
Administer the below before rituximab :				
<input checked="" type="checkbox"/> acetaminophen	650 mg, Tab, PO, Once, Comment: to be given prior to rituximab infusion			
<input checked="" type="checkbox"/> diphenhydrAMINE	25 mg, Injection, IV Push, Once, Comment: to be given prior to rituximab infusion			
CHEMOTHERAPY				
	Drug (generic) & solution (optional)	Intended Dose	Actual Dose	Route, Infusion, Frequency and total doses
<input checked="" type="checkbox"/>	rituximab	375 mg/m²		IV Piggyback, Infuse using Rituximab flowsheet, ONCE on DAY 1
<input checked="" type="checkbox"/>	fludarabine	25 mg/m²		IV Piggyback, Infuse over 30 min, q24h on DAYS 2,3, and 4
<input checked="" type="checkbox"/>	cyclophosphamide	250 mg/m²		IV Piggyback, Infuse over 2 hours, q24h on DAY 2, 3, and 4
Acute Emesis Prophylaxis (may undergo therapeutic interchange)				
NOTE: Administer intial doses at least 30-60 minutes prior to chemotherapy				
<input checked="" type="checkbox"/>	ondansetron	12 mg, Injection, IV Piggyback, Once, DAY 2-4		
<input checked="" type="checkbox"/>	prochlorperazine	10 mg, Injection, IV Push, q6h, PRN Nausea/Vomiting , Comment : if unable to take PO		
<input checked="" type="checkbox"/>	prochlorperazine	10 mg, Tab, PO, q6h, PRN Nausea/Vomiting		
Consults/Notifications				
<input type="checkbox"/>	Notify Physician- Once	T;N, Who: _____, For: if BSA exceeds 2 m ²		

Date

Time

Physician's Signature

MD Number