



Physician Orders ADULT: Tolvaptan and Conivaptan Plan

Tolvaptan and Conivaptan Plan

Vital Signs

- Vital Signs Per Unit Protocol
T;N, Routine Monitor and Record T,P,R,BP, q6h(std)

Patient Care

- Intake and Output
T;N, Routine, q8h(std)
- INT Insert/Site Care
T;N, Routine, QDay, Inspect conivaptan IV site for phlebitis
- INT Insert/Site Care
T;N, QDay, Rotate IV site of conivaptan
- Tolvaptan/Conivaptan Ordering Criteria

Nursing Communication

- Nursing Communication
T;N, If Na level rise is equal to or greater than 10 mEq over 24 hours, notify Physician

Continuous Infusion

- conivaptan
20 mg, IV Piggyback, IV Piggyback, q24h, (for 96 hr), (infuse over 24 hr)
Comments: give after IV bolus. rotate IV site every 24 hours unless infused centrally. 96 hour stop

Medications

- CHOOSE ONE OF THE MEDICATIONS BELOW(NOTE)*
- +1 Hours** conivaptan
20 mg, IV Piggyback, IV Piggyback, once, (infuse over 30 min)
Comments: give before continuous infusion. Rotate IV site every 24 hours unless infused centrally.
 - +1 Hours** tolvaptan
 - 15 mg, Tab, PO, QDay, (for 4 dose) (DEF)**
 - 30 mg, Tab, PO, QDay, (for 4 dose)*

Laboratory

- Na Level
Time Study, q6h x 24 hr, Type: Blood
- Na Level
Routine, T;N+2880, QDay, Type: Blood

Consults/Notifications/Referrals

- Notify Physician-Continuing
T;N, if suspected phlebitis at IV site
- Pharmacy Consult - Tolvaptan/Conivaptan Monitoring
T;N, Routine, QDay



Attach patient label here



Physician Orders ADULT: Tolvaptan and Conivaptan Plan

Date Time Physician's Signature MD Number

***Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

