



Physician Orders PEDIATRIC: LEB GEN SURG Pectus Excav POD 1 Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- Initiate Powerplan Phase*
Phase: LEB GEN SURG Pectus Excav POD1 Phase, When to Initiate: _____

LEB GEN SURG Pectus Excav POD 1 Phase

Activity

- Out Of Bed
Up To Chair, bid, with PT assistance for the first time out of bed
- Ambulate
bid, With Assistance, with PT assistance for the first time ambulating

Food/Nutrition

- Regular Pediatric Diet

Patient Care

- DC IV Fluids When Tolerating PO
T;N, DC IV Fluids and Heplock IV when tolerating PO
- Foley Remove
- Indwelling Urinary Catheter Insert-Follow Removal Protocol

Nursing Communication

- Nursing Communication
Please discontinue PCA basal rate.

Continuous Infusion

- D5 1/2NS
1,000 mL, IV, Routine, TKO
Comments: To infuse TKO with PCA .

Medications

- +1 Hours** bisacodyl
10 mg, Supp, PR, QDay, Routine, May hold if patient has had a BM in last 24 hours
- +1 Hours** docusate
100 mg, Cap, PO, bid, Routine
- +1 Hours** methocarbamol
500 mg, Tab, PO, tid, Routine
- +1 Hours** acetaminophen-oxyCODONE 325 mg-5 mg oral tablet
1 tab, Tab, PO, q4h, PRN Pain, Moderate (4-7), Routine, (for 3 day), (1 tab = 5mg oxyCODONE)

Consults/Notifications/Referrals

- Notify Physician-Continuing
Notify: physician, Notify For: if unable to void 8 hours after Foley removal
- Pharmacy Consult
Reason: Discontinue PCA basal rate

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

***Report Legend:**
 DEF - This order sentence is the default for the selected order
 GOAL - This component is a goal
 IND - This component is an indicator
 INT - This component is an intervention
 IVS - This component is an IV Set
 NOTE - This component is a note
 Rx - This component is a prescription
 SUB - This component is a sub phase, see separate sheet

Attach patient label here



Physician Orders PEDIATRIC: LEB GEN SURG Pectus Excav POD 1 Plan

R-Required order

