

	Orders Phase ets/Protocols/PowerPlans
	Initiate Powerplan Phase Phase: Heart Failure Admit Phase, When to Initiate:
Heart F	Failure Admit Plan
	ategorized
R	Heart Failure Care Track
R_	Heart Failure Quality Measures
	Add To Problem List
_	Problem: Heart failure
Ш	Add To Problem List
_	sion/Transfer/Discharge
	Patient Status Initial Inpatient
	T;N Admitting Physician:
	Bed Type: Specific Unit:
	Care Team: Anticipated LOS: 2 midnights or more
	Patient Status Initial Outpatient
	T;N, Attending Physician:
	Reason for Visit:
	Bed Type:Specific Unit:
	Outpatient Status/Service OP-OBSERVATION Services
	Notify Dhysician Onco
	Notify Physician-Once Notify For: room number upon admission
Vital S	
$\overline{\mathbf{v}}$	Vital Signs
	Monitor and Record T,P,R,BP, q4h(std) x 24hr, then q8h(std)
$\overline{\mathbf{A}}$	Orthostatic Blood Pressure
	QDay, and orthostatic pulse rate
Activit	y
	Bedrest
	Bedrest w/BRP
$\overline{\mathbf{A}}$	Ambulate
	tid, ad lib
	lutrition
	Low Sodium Diet
	American Heart Association Diet
Ш	Consistent Carbohydrate Diet
	Start at: T;N, 1200 Calorie, Insulin: [] No Insulin [] Short Acting
	[] Intermediate [] Long Acting [] Short and Intermediate [] Short and Long;
	Renal Patient:[] No [] Yes, on dialysis [] Yes, not on dialysis
	Start at: T;N, Caloric Level: 1800 Calorie, Insulin: [] No Insulin [] Short Acting
	[] Intermediate [] Long Acting [] Short and Intermediate [] Short and Long; Renal Patient:[] No [] Yes, on dialysis [] Yes, not on dialysis
	□ Start at: T;N, 2400 Calorie, Insulin: [] No Insulin [] Short Acting [] Intermediate [] Long Acting [] Short and Intermediate [] Short and Long;
	Renal Patient:[] No [] Yes, on dialysis [] Yes, not on dialysis
Patient	
	Code Status
	Resus Type: DNR-Do Not Resuscitate





$\overline{\mathbf{A}}$	Intake and Output q2h x 3 occurrences, then q6h
$\overline{\mathbf{Z}}$	Urine Output Goal Goal: >500mL / 6 hrs
$\overline{\mathbf{v}}$	Daily Weights
	qam IV Insert/Site Care
	q4day, establish IV access Restrict Fluids
R	1000 mL/ 24 h Smoking Cessation Advice/Counseling
R —	Instruct/Educate patient and family on CHF and give CHF materials and document.
☐ Nursin	Bedside Glucose Nsg g Communication
$\overline{\mathbf{A}}$	Nursing Communication if urine output > 1000 mL/6 hours, order STAT BMP and magnesium level
$\overline{\mathbf{v}}$	Nursing Communication
	if patient has a foley catheter, remove it when diuretic is PO and patient is able to void and save urine independently
☑	Nursing Communication retrieve previous ECHO, cath, or MUGA report from Cerner and document ejection fraction result, and date of the report and if study showed LVSD.
☑	Nursing Communication Consult company representative to evaluate need for wearable cardiac defibrillator for new onset heart failure with Left Ventricular Ejection Fraction less than 35%
Respir	atory Care
☑	O2-Nasal Cannula 2 L/min, Special Instructions: titrate to keep o2 sat =/>92%
	O2 Sat-Spot Check (RT) QDay, Special Instructions: room air
Medica	
$\overline{\mathbf{C}}$	VTE MEDICAL Prophylaxis Plan(SUB)*
	+1 Hours furosemide
	40 mg, Injection, IV Push, q8h, Routine
	Alternate Dose Below(NOTE)*
	+1 Hours furosemide
	40 mg, Injection, IV Push, q12h, Routine To adhere to Regulatory guidelines, if ACEI/ARB therapy is contraindicated and will not be ordered as a
	discharge prescription, document the Reason ACEI Not Prescribed at Discharge AND the Reason ARB Not Prescribed at Discharge below:(NOTE)*
	+1 Hours lisinopril 10 mg, Tab, PO, QDay, Routine
	+1 Hours quinapril
	5 mg, Tab, PO, QDay, Routine +1 Hours benazepril
_	10 mg, Tab, PO, QDay, Routine
	+1 Hours hydrALAZINE 35 mg, Tab, PO, q8h, Routine
	+1 Hours isosorbide dinitrate 20 mg, Tab, PO, q8h, Routine





	Reason ACEI Not Prescribed at Discharge
	Reason ARB Not Prescribed at Discharge
	Nitrate Medication Orders (see below)(NOTE)*
	Nursing Communication
	Hold nitroglycerin if SBP less than 100mmHg
	+1 Hours nitroglycerin
	0.4 mg, Tab, SL, q5min, PRN Chest Pain, STAT, (for 3 occurrence)
_	Comments: (3 occurence)
	+1 Hours metoprolol extended release
_	12.5 mg, ER Tablet, PO, QDay, Routine
	+1 Hours carvedilol
_	3.125 mg, Tab, PO, bid, Routine
	+1 Hours spironolactone
_	25 mg, Tab, PO, QDay, Routine
	PRN Common Comfort Medication Plan(SUB)*
	+1 Hours acetaminophen
_	650 mg, Tab, PO, q6h, PRN Pain, Mild or Fever, Routine
	+1 Hours promethazine
_	12.5 mg, Tab, PO, q4h, PRN Nausea/Vomiting, Routine
	+1 Hours ondansetron
_	4 mg, Injection, IV Push, q6h, PRN Nausea/Vomiting, Routine
	+1 Hours temazepam
	15 mg, Cap, PO, hs, PRN Insomnia, Routine
	+1 Hours famotidine
	20 mg, Tab, PO, q12h, PRN Indigestion, Routine
	Comments: reduce to 20 mg daily if creatinine clearance is less than 50 mL/min
ш	+1 Hours famotidine
	20 mg, Injection, IV Push, q12h, PRN Indigestion, Routine Comments: reduce to 20 mg daily if creatinine clearance is less than 50 mL/min
	+1 Hours magnesium hydroxide
_	30 mL, Oral Susp, PO, QDay, PRN Constipation, Routine
	+1 Hours bisacodyl
_	10 mg, DR Tablet, PO, once
	+1 Hours Maalox Advanced Maximum Strength
_	10 mL, Oral Susp, PO, q6h, PRN Indigestion, Routine
	Insulin SENSITIVE Sliding Scale Plan(SUB)*
	Insulin STANDARD Sliding Scale Plan(SUB)*
	Insulin RESISTANT Sliding Scale Plan(SUB)*
_ _abora	
	Troponin-I
_	Time Study, T;N, once, Type: Blood
$\overline{\mathbf{v}}$	Troponin-I
_	Time Study, T;N+180, once, Type: Blood
$\overline{\mathbf{A}}$	Troponin-I
_	Time Study, T;N+360, once, Type: Blood
$\overline{\checkmark}$	CMP
_	Routine, T;N, once, Type: Blood
$\overline{\checkmark}$	Magnesium Level
	Routine, T;N, once, Type: Blood





$\overline{\mathbf{A}}$	BNP Pro
	Routine, T;N, once, Type: Blood Brain Natriuretic Peptide
V	Routine, T;N, once, Type: Blood CBC w/o Diff
_	Routine, T;N, once, Type: Blood
$\overline{\mathbf{C}}$	Urinalysis w/Reflex Microscopic Exam
$\overline{\mathbf{Q}}$	Routine, T;N, once, Type: Urine, Nurse Collect
	Lipid Profile Routine, T+1;0400, once, Type: Blood Comments: fasting
$\overline{\mathbf{v}}$	BMP
	Routine, T+1;0400, qam x 3 day, Type: Blood
Diagno	ostic Tests
	EKG
_	Start at: T;N, Priority: Routine, Reason: Congestive Heart Failure, Frequency: once
	Chest 2 Views T;N, Reason for Exam: Congestive Heart Failure, Routine, Stretcher
	Chest 1 View
	T;N, Reason for Exam: Congestive Heart Failure, Routine, Portable
	TTE Echo W/Contrst or 3D if needed
	lts/Notifications/Referrals
☑	Notify Physician-Continuing Notify For: if potassium < 4mmol/L, magnesium < 2mmol/L, or urine output goal is not met
$\overline{\checkmark}$	Consult Clinical Dietitian
	Type of Consult: Other, please specify, Special Instructions: low sodium diet counseling
☑	Cardiac Rehab Consult/Doctor Order T;N
	Comments: Walk patient in hallway, CHF education.
	Discharge Planning
	Refer patient to Methodist heart failure clinic: 516-2442
	Consult Case Management
	Consult Medical Social Work
Date	Time Physician Signature MD Number

*Report Legend:

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Require order

