



attach patient label

Physician Orders

LEB Med-Surg Intranasal Medications Plan

[X or R] = will be ordered unless marked out.

PEDIATRIC

Height: _____ cm Weight: _____ kg

Allergies: No known allergies

Medication allergy(s): _____

Latex allergy Other: _____

Patient Care

O2 Sat Monitoring NSG T;N, Place on patient prior to administration of Intranasal medication and continue monitoring until 30 mins following procedure completion.

Nursing Communication

Nursing Communication T;N, Once procedure completed, nurse to discontinue LEB Med-Surg Intranasal Medications Plan

Medication

midazolam 0.2 mg/kg, Inh Soln, Nasal, N/A, PRN Other:Specify in Comment, Anxiolysis, STAT, Max Dose = 10 mg; may repeat dose x one dose 15 mins after 1st dose per MD discretion.

midazolam 0.3 mg/kg, Inh Soln, Nasal, N/A, PRN Other:Specify in Comment, Anxiolysis, STAT, Max Dose = 10 mg; may repeat dose x one dose 15 mins after 1st dose per MD discretion.

fentaNYL 2 mcg/kg, Inh Soln, Nasal, N/A, PRN Pain, STAT, Max Dose = 100 mcg; may repeat dose once, may repeat dose x one dose 15 mins after 1st dose per MD discretion.

fentaNYL 3 mcg/kg, Inh Soln, Nasal, N/A, PRN Pain, STAT, Max Dose = 100 mcg; may repeat dose once, may repeat dose x one dose 15 mins after 1st dose per MD discretion.

NOTE: Order buffered lidocaine in conjunction with midazolam to prevent nasal irritation.

buffered lidocaine 1% inj 1 mL, Inh Soln, Nasal, once, STAT, 0.5mL in each nostril 2-3 minutes prior to intranasal midazolam to prevent nasal burning

NOTE: Order Naloxone if fentaNYL is ordered

naloxone 0.02mg/kg, Ped Injectable, IM, N/A, STAT, Opioid Reversal. Min. dose = 0.1mg. May repeat up to max dose - 2mg

naloxone 0.02mg/kg, Ped Injectable, IV Push, N/A, STAT, Opioid Reversal. Min. dose = 0.1mg. May repeat up to max dose - 2mg

Date Time Physician's Signature MD Number

