



## Physician Orders ADULT: PCA - MorPHINE Protocol Plan (Adult)

### PCA - MorPHINE Protocol Plan (Adult)

#### Vital Signs

R Vital Signs

*Routine Monitor and Record Pulse | Resp Rate | Blood Pressure, Quality of Respirations, Oxygen Saturation; q30min x2, q1h x1, then q4h until PCA is Discontinued per hospital policy., T;N*

#### Patient Care

☒ Instruct/Educate

*Instruct: patient and family, Method: Demonstrate, Topic: proper administration and use of PCA*

R Pain Score

*MorPHINE PCA Protocol Plan(Adult): q30min x2; q1h x1; then q4h per hospital policy. T;N*

#### Nursing Communication

R Nursing Communication

*MorPHINE PCA Protocol Plan(Adult): If no IV Fluids are ordered, enter an order for NS to infuse at TKO rate while on PCA therapy. T;N*

R Nursing Communication

*MorPHINE PCA Protocol Plan(Adult): Place pulse oximeter on patient for oxygen saturation monitoring and documentation, T;N*

R Nursing Communication

*MorPHINE PCA Protocol Plan(Adult): PCA Titration Band, document once per shift via Iview>>Blood Admin and Titrations>>IV drips>> locate section for PCA. Document all components; DO NOT CLEAR the PCA until after you have documented this information. T;N*

#### Medications

Changes to the protocol are not permitted if you deviate from the protocol you must specify all parameters of PCA Orders (concentration, loading dose, bolus dose, lockout, basal rate and 1 hour limit.(NOTE)\*

Use standard concentration PCA medications for opioid naive patients and patients with normal dosing requirements.(NOTE)\*

For patients with severe pain, who have very high PCA requirements (greater than or equal to morPHINE 50 mg IV per shift), order the "Maximally Concentrated PCA Orders for Adults" instead.(NOTE)\*

To order "per protocol" select one of the orders below and leave unchanged. The first order is without a basal rate. The second order includes a standard basal rate.(NOTE)\*

Choose one of the following orders below:(NOTE)\*

☒ **+1 Hours** MorPHINE PCA - 1 mg/mL

*PCA Dose: 1 mg, Lock-out: 10 min, 1hr Limit: 6 mg, Basal Rate: No Basal Rate, PCA, PCA, Routine*

☐ **+1 Hours** MorPHINE PCA - 1 mg/mL

*PCA Dose: 1 mg, Lock-out: 10 min, 1hr Limit: 7 mg, Basal Rate: 1 mg/hr, PCA, PCA*

Note: All active opioid orders will be discontinued by the reviewing pharmacist upon receiving an order for a PCA. Orders for breakthrough pain should be ordered with PCA.(NOTE)\*

R naloxone

*0.4 mg, Injection, IV Push, q5min, PRN Respiratory Depression, Routine  
Comments: For Respiratory Rate less than 12.*

☐ ondansetron

*4 mg, Orally Disintegrating Tab, PO, q6h, PRN Nausea/Vomiting, Routine*

☐ ondansetron

*4 mg, Injection, IV Push, q6h, PRN Nausea/Vomiting, Routine  
Comments: May have IV if unable to tolerate PO*

☐ diphenhydramine

*25 mg, Cap, PO, q4h, PRN Itching, Routine*

☐ diphenhydramine

*12.5 mg, Injection, IV Push, q4h, PRN Itching, Routine  
Comments: May have IV if unable to tolerate PO*

☐ docusate

*100 mg, Cap, PO, bid, Routine*

☐ senna

*8.6 mg, Tab, PO, bid, PRN Constipation, Routine*





**Physician Orders ADULT: PCA - MorPHINE Protocol Plan (Adult)**

*Comments: Once tolerating diet or PO meds*

- ☐ bisacodyl  
10 mg, Tab, PO, QDay, PRN Constipation, Routine
- ☒ polyethylene glycol 3350  
17 g, Powder, PO, q24h, PRN Constipation, Routine

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

**\*Report Legend:**

DEF - This order sentence is the default for the selected order  
GOAL - This component is a goal  
IND - This component is an indicator  
INT - This component is an intervention  
IVS - This component is an IV Set  
NOTE - This component is a note  
Rx - This component is a prescription  
SUB - This component is a sub phase, see separate sheet  
R-Required

