



**Physician Orders PEDIATRIC: LEB Cardiovascular Surgery Transfer Plan**

**Initiate Orders Phase**

**Care Sets/Protocols/PowerPlans**

- Initiate Powerplan Phase  
*Phase: LEB Cardiovascular Surgery Transfer Phase, When to Initiate: When patient arrives to unit*

**LEB Cardiovascular Surgery Transfer Phase**

**Admission/Transfer/Discharge**

- Transfer Pt within current facility
- Notify Physician-Once  
*Notify For: of room number on arrival to unit*

**Vital Signs**

- Vital Signs
  - Monitor and Record T,P,R,BP, q4h(std), omit 0200 VS (DEF)\*
  - Monitor and Record T,P,R,BP, q1h(std)
  - Monitor and Record T,P,R,BP, q2h(std)
  - Monitor and Record T,P,R,BP, q4h(std)

**Activity**

- Activity As Tolerated
- Bedrest

**Food/Nutrition**

- NPO
- Breastfeed  
*T;N*
- LEB Formula Orders Plan(SUB)\*
- Clear Liquid Diet  
*Start at: T;N*
- Low Fat Diet
- No Fat Diet <10g per day
- Regular Pediatric Diet

**Patient Care**

- Advance Diet As Tolerated  
*Start with clear liquid diet and advance as tolerated*
- Daily Weights  
*qEve*
- O2 Sat Monitoring NSG  
*Routine*
- Cardiopulmonary Monitor  
*Monitor Type: CP Monitor*
- Telemetry
  - Routine (DEF)\*
  - Routine, May be off monitor to walk in hall or for xray and ECHO done off unit
- Elevate Head Of Bed  
*If patient is 18 years old or greater place both orders below for VTE Prophylaxis:(NOTE)\**
- Consult Clinical Pharmacist  
*Reason: assess for VTE Prophylaxis*
- SCD Apply
- Incentive Spirometry NSG  
*q1h-Awake*

**Respiratory Care**





### Physician Orders PEDIATRIC: LEB Cardiovascular Surgery Transfer Plan

- LEB NICU Respiratory Plan(SUB)\*  
 Oxygen Delivery  
*Special Instructions: Titrate to keep O2 sat at \_\_\_ to \_\_\_%*

#### Medications

- +1 Hours** diphenhydrAMINE  
 1 mg/kg, *Ped Injectable, IV, q8h, PRN Other, specify in Comment, Routine (DEF)\**  
*Comments: Agitation , Max dose = 50mg*  
 12.5 mg, *Ped Injectable, IV, q8h, PRN Other, specify in Comment, Routine*  
*Comments: Agitation*  
 25 mg, *Ped Injectable, IV, q8h, PRN Other, specify in Comment, Routine*  
*Comments: Agitation*
- +1 Hours** ondansetron  
 0.1 mg/kg, *Oral Susp, PO, q8h, PRN Nausea/Vomiting, Routine, Max dose = 8 mg (DEF)\**  
 4 mg, *Orally Disintegrating Tab, PO, q8h, PRN Nausea/Vomiting, Routine, (for 3 dose )*
- +1 Hours** ondansetron  
 0.1 mg/kg, *Ped Injectable, IV Piggyback, q6h, PRN Nausea/Vomiting, Routine, Max dose = 8 mg*  
*Comments: May give IV if unable to take PO*
- +1 Hours** famotidine  
 0.25 mg/kg, *Injection, IV, q12h, Routine, Max Daily Dose = 40 mg/day*

#### Analgesics

- +1 Hours** ibuprofen  
 200 mg, *Tab, PO, q8h, PRN Pain, Mild (1-3), Routine (DEF)\**  
 10 mg/kg, *Tab, PO, q8h, PRN Pain, Mild (1-3), Routine*  
 10 mg/kg, *Oral Soln, PO, q8h, PRN Pain, Mild (1-3), Routine*
- +1 Hours** acetaminophen-HYDROcodone 325 mg-5 mg oral tablet  
 1 tab, *Tab, PO, q4h, PRN Pain, Moderate (4-7), Routine, (for 5 day ), (1 tab = 5 mg HYDROcodone)*
- +1 Hours** acetaminophen-HYDROcodone 325 mg-7.5 mg/15 mL oral solution  
 0.15 mg/kg, *Elixir, PO, q6h, PRN Pain, Moderate (4-7), Routine, (for 5 day ), (5 mL = 2.5 mg HYDROcodone), Max dose = 10mg*
- +1 Hours** acetaminophen-HYDROcodone 325 mg-7.5 mg oral tablet  
 1 tab, *Tab, PO, q4h, PRN Pain, Moderate (4-7), Routine, (for 5 day ), (1 tab = 7.5 mg of Hydrocodone)*

#### Anti-pyretics

- +1 Hours** acetaminophen  
 10 mg/kg, *Liq, PO, q4h, PRN Fever, Routine, For temperature greater than 38, Max Dose = 75 mg/kg/day up to 4g/day (DEF)\**  
 325 mg, *Tab, PO, q4h, PRN Fever, Routine, For temperature greater than 38, Max Dose = 75 mg/kg/day up to 4g/day*
- +1 Hours** acetaminophen  
 10 mg/kg, *Supp, PR, q4h, PRN Fever, Routine, For temperature greater than 38, Max Dose = 75 mg/kg/day up to 4g/day*  
*Comments: May give PR if unable to take PO*

#### Bowel Care

- +1 Hours** glycerin pediatric rectal suppository  
 1 supp, *Supp, PR, q24h, PRN Other, specify in Comment, Routine, For no stool for 24 hours*
- +1 Hours** polyethylene glycol 3350  
 8.5 g, *Powder, PO, QDay, PRN Constipation, Routine (DEF)\**  
 17 g, *Powder, PO, QDay, PRN Constipation, Routine*





### Physician Orders PEDIATRIC: LEB Cardiovascular Surgery Transfer Plan

- +1 Hours** docusate
- 2.5 mg/kg, Oral Susp, PO, bid, Routine, Max dose = 100mg (1 mL = 10 mg) (DEF)\*
  - 50 mg, Cap, PO, bid, Routine
  - 100 mg, Cap, PO, bid, Routine

#### Diuretics

- +1 Hours** furosemide
- 1 mg/kg, Oral Soln, PO, q8h, Routine, Max dose = 40 mg (5 mL = 50 mg) (DEF)\*
  - 1 mg/kg, Tab, PO, q8h, Routine, Max dose = 40 mg
  - 1 mg/kg, Oral Soln, PO, q12h, Routine, Max dose = 40 mg (5 mL = 50 mg)
  - 1 mg/kg, Tab, PO, q12h, Routine, Max dose = 40 mg
- +1 Hours** furosemide
- 1 mg/kg, Ped Injectable, IV Push, q8h, Routine, Max dose = 40 mg (DEF)\*
  - 1 mg/kg, Ped Injectable, IV Push, q12h, Routine, Max dose = 40 mg

#### Respiratory Medications

- +1 Hours** albuterol (MDI)
- 2 puff, MDI, INH, q4h, Routine, (for 6 dose ), (1 puff = 90 mcg)

#### Laboratory

- CBC  
*Routine, T;N, once, Type: Blood*
- CBC  
*Routine, T+1;0400, once, Type: Blood*
- BMP  
*Routine, T;N, once, Type: Blood*
- BMP  
*Routine, T+1;0400, once, Type: Blood*
- PT/INR  
*Routine, T;N, once, Type: Blood*
- PT/INR  
*Routine, T+1;0400, once, Type: Blood*
- PTT  
*Routine, T;N, once, Type: Blood*
- PTT  
*Routine, T+1;0400, once, Type: Blood*

#### Diagnostic Tests

- Chest 1 View  
*T+1;0800, Reason for Exam: Other, Enter in Comments, Routine, Portable*  
*Comments: Postop cardiovascular surgery*
- Echocardiogram Pediatric (0-18 yrs)  
*Start at: T;N, Priority: Routine, Reason: Other, specify, Other reason: postop cardiovascular surgery,*  
*Transport: Bedside*  
*Comments: Other reason: postop cardiovascular surgery*

#### Consults/Notifications/Referrals

- Notify Physician For Vital Signs Of  
*Notify: Cardiology/Cardiology Fellow*
- Notify Physician For Vital Signs Of  
*Notify: CV Surgeon*
- Notify Physician-Continuing
- Consult MD Group





**Physician Orders PEDIATRIC: LEB Cardiovascular Surgery Transfer Plan**

- Consult MD
- Dietitian Consult/Nutrition Therapy
- Lactation Consult  
*Routine*
- Medical Social Work Consult  
*Reason: Assistance at Discharge*

---

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

**\*Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

