



Physician Orders PEDIATRIC: LEB UROL General Admit Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- Initiate Powerplan Phase
Phase: LEB UROL General Admit Phase, When to Initiate: _____

LEB UROL General Admit Phase

Admission/Transfer/Discharge

- Patient Status Initial Outpatient
T;N Attending Physician: _____
Reason for Visit: _____
Bed Type: _____ Specific Unit: _____
Outpatient Status/Service: [] Ambulatory Surgery, [] OP Diagnostic Procedure
[] OP OBSERVATION Services

Vital Signs

- Vital Signs
 - Monitor and Record T,P,R,BP, q4h(std) (DEF)*
 - Monitor and Record T,P,R,BP, post op, then q4h

Activity

- Bedrest
Routine
- Activity As Tolerated
Up Ad Lib
- Out Of Bed
tid

Food/Nutrition

- NPO Communication Nsg
After Midnight
- NPO
- Breastfeed
- LEB Formula Orders Plan (SUB)*
- Regular Pediatric Diet
- Clear Liquid Diet
Start at: T;N

Patient Care

- Advance Diet As Tolerated
Start clear liquids and advance to regular diet as tolerated.
- Intake and Output
Routine, q2h (std)
- Foley Care
to gravity
- Dressing Care
- Drain Care
- Supply Request CSR
Geomat
- Cardiopulmonary Monitor
Routine, Monitor Type: CP Monitor
- Incentive Spirometry NSG
q1h-Awake

Respiratory Care

- RT Assess and Call





Physician Orders PEDIATRIC: LEB UROL General Admit Plan

Routine, Special Instructions: Bronchial Hygiene (BHH) Protocol

Continuous Infusion

- D5 1/2NS
1,000 mL, IV, Routine, mL/hr
- D5 1/4 NS
1,000 mL, IV, Routine, mL/hr
- Sodium Chloride 0.9%
1,000 mL, IV, Routine, mL/hr
- 1/2NS
1,000 mL, IV, Routine, mL/hr

Medications

- +1 Hours** diphenhydrAMINE
1 mg/kg, Elixir, PO, q4h, PRN Itching, Routine, max dose = 50mg, (5mL = 12.5mg)
- +1 Hours** diphenhydrAMINE
1 mg/kg, Ped Injectable, IV, q4h, PRN Itching, Routine, max dose 50 mg
- +1 Hours** B & O Suppettes 15-A
 - 0.25 supp, Supp, PR, q6h, PRN Bladder Spasm, Routine (DEF)*
 - 0.33 supp, Supp, PR, q6h, PRN Bladder Spasm, Routine
 - 0.5 supp, Supp, PR, q6h, PRN Bladder Spasm, Routine
 - 1 supp, Supp, PR, q6h, PRN Bladder Spasm, Routine
- +1 Hours** hyoscyamine elixir
 - 31.25 mcg, Elixir, PO, q4h, PRN Bladder Spasm, Routine, (1.25 mL = 31.25 mcg) (DEF)*
 - 62.5 mcg, Elixir, PO, q4h, PRN Bladder Spasm, Routine, (2.5 mL = 62.5 mcg)
- +1 Hours** hyoscyamine
0.125 mg, Tab, PO, q4h, PRN Bladder Spasm, Routine
- +1 Hours** oxybutynin
 - 0.2 mg/kg, Susp, PO, tid, Bladder Spasm, Routine, 1 to 5 years (DEF)*
 - 0.2 mg/kg, Tab, PO, tid, Bladder Spasm, Routine, 1 to 5 years
- +1 Hours** oxybutynin extended release
5 mg, ER Tablet, PO, QDay, Routine, greater than or equal to 6 years
- +1 Hours** Flomax
0.2 mg, Cap, PO, hs, Routine
- +1 Hours** furosemide
1 mg/kg, Tab, PO, q6h, Routine
- +1 Hours** furosemide
1 mg/kg, Ped Injectable, IV Push, q6h, Routine

Anti-infectives

- +1 Hours** Triple Antibiotic
1 application, Ointment, TOP, tid, PRN Wound Care, Routine, apply to stoma side bid-tid
- +1 Hours** ampicillin
25 mg/kg, Ped Injectable, IV Piggyback, q6h, Routine, (for 14 day), Max dose 12 grams/day
- +1 Hours** cefTRIAxone
50 mg/kg, Ped Injectable, IV Piggyback, q24h, Routine, (for 14 day), Max dose = 2 grams
- +1 Hours** clindamycin
 - 10 mg/kg, Oral Soln, PO, q8h, Routine, (for 14 day) (DEF)*
 - 10 mg/kg, Cap, PO, q8h, Routine, (for 14 day)
- +1 Hours** clindamycin
10 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 14 day)





Physician Orders PEDIATRIC: LEB UROL General Admit Plan

- +1 Hours erythromycin base**
 - 20 mg/kg, Oral Susp, PO, tid, Routine, (for 3 dose), To be given at 1300, 1400 and 2300, Bowel Prep (DEF)*
 - 20 mg/kg, Tab, PO, tid, Routine, (for 3 dose), To be given at 1300, 1400 and 2300, Bowel Prep
- +1 Hours neomycin**
25 mg/kg, Tab, PO, q8h, Routine, (for 3 dose), To be given at 1300, 1400 and 2300, Bowel Prep
- +1 Hours ceFAZolin**
25 mg/kg, Ped Injectable, IV Piggyback, q8h, Routine, (for 14 day)
- +1 Hours Gentamicin Bladder Irrigation 0.48 mg/mL (Pediatric)**
30 mL, Topical Soln, IRR, QNight, Routine
Comments: Instill into bladder
- +1 Hours cephalixin**
12.5 mg/kg, Oral Susp, PO, q6h, Routine, (for 14 day), Max dose = 500 mg
- +1 Hours metroNIDAZOLE**
7.5 mg/kg, Ped Injectable, IV Piggyback, q6h, Routine, (for 14 day), Max dose = 4grams/day,
- +1 Hours nitrofurantoin**
 - 2 mg/kg, Oral Soln, PO, QDay, Routine, (for 14 day), Max dose = 100 mg/day, UTI Prophylaxis (DEF)*
 - 50 mg, Cap, PO, QDay, Routine, (for 14 day), UTI Prophylaxis
 - 100 mg, Cap, PO, QDay, Routine, (for 14 day), UTI Prophylaxis
- +1 Hours sulfamethoxazole-trimethoprim susp**
2 mg/kg, Susp, PO, q24h, Routine, (for 14 day), UTI Prophylaxis, dosed as mg of TMP

Analgesics

- +1 Hours acetaminophen**
 - 10 mg/kg, Liq, PO, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day (DEF)*
 - 80 mg, Chew tab, PO, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day
 - 325 mg, Tab, PO, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day
- +1 Hours acetaminophen**
10 mg/kg, Supp, PR, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day
Comments: May give suppository if unable to take oral medication.
- +1 Hours acetaminophen-HYDROcodone 325 mg-7.5 mg/15 mL oral solution**
0.15 mg/kg, Elixir, PO, q6h, PRN Pain, Moderate (4-7), Routine, (for 5 day), (5 mL = 2.5 mg HYDROcodone), Max dose = 10mg
- +1 Hours acetaminophen-HYDROcodone 325 mg-5 mg oral tablet**
1 tab, Tab, PO, q4h, PRN Pain, Moderate (4-7), Routine, (for 5 day), (1 tab = 5 mg HYDROcodone), Max dose = 10 mg
- +1 Hours morphine**
0.1 mg/kg, Ped Injectable, IV, q3h, PRN Pain, Breakthrough, Routine, (for 3 day), Max initial dose = 2mg

Bowel Preparation

- +1 Hours bisacodyl**
10 mg, Supp, PR, QDay, PRN Constipation, Routine
- +1 Hours polyethylene glycol 3350**
1 g/kg, Powder, PO, QDay, Routine
- +1 Hours magnesium citrate**
 - 3 mL/kg, Liq, PO, q6h, Routine, (for 2 dose) [Less Than 7 year] (DEF)*
 - 100 mL, Liq, PO, q6h, Routine, (for 2 dose) [7 - 12 year]





Physician Orders PEDIATRIC: LEB UROL General Admit Plan

- 150 mL, Liq, PO, q6h, Routine, (for 2 dose) [Greater Than or Equal To 12 year]
- +1 Hours** sodium biphosphate-sodium phosphate enema pediatric
66 mL, Enema, PR, once, Routine, (2 to 11 years)
- +1 Hours** sodium biphosphate-sodium phosphate enema adult
133 mL, Enema, PR, once, Routine

Antiemetics

- +1 Hours** ondansetron
 - 0.1 mg/kg, Oral Susp, PO, q8h, PRN Nausea/Vomiting, Routine, Max dose = 4mg (DEF)*
 - 4 mg, Orally Disintegrating Tab, PO, q8h, PRN Nausea/Vomiting, Routine, dose)
- +1 Hours** ondansetron
0.1 mg/kg, Ped Injectable, IV Push, q8h, PRN Nausea/Vomiting, Routine, Max dose = 8 mg

Laboratory

- CBC
T;N, Routine, once, Type: Blood
- BMP
T;N, Routine, once, Type: Blood
- Urinalysis w/Reflex Microscopic Exam
Routine, T;N, once, Type: Urine
- Urine Culture
Routine, T;N, Specimen Source: Urine

Diagnostic Tests

- Abdomen 1 View
T;N, Routine, Wheelchair
- LEB CT Abdomen W Cont Plan (SUB)*
- LEB CT Abdomen WWO Cont Plan (SUB)*
- LEB CT Pelvis W Cont Plan (SUB)*
- LEB CT Pelvis WO Cont Plan (SUB)*
- Pyelogram IV WWO KUB WWO Tomography
T;N, Routine, Wheelchair
- Urethrocytogram Retrograde
T;N, Routine, Wheelchair
- LEB US Abd Comp w/Delay Diet Plan (SUB)*
- US Retroperitoneal B Scan/Real Time Comp
T;N, Routine, Wheelchair
- NM Kidney Vasc Flow & Funct Sing W Phar
T;N, Routine, Wheelchair
- NM Kidney Imaging Morphology
T;N, Routine, Wheelchair
- Urodynamics Evaluation LEB

Consults/Notifications/Referrals

- Notify Physician-Once
Notify For: of room number on arrival to unit
- Notify Physician-Continuing
Notify: Urology oncall for poor tolerance of bowel prep or questions
- Notify Physician For Vital Signs Of
- Consult MD Group
- Consult MD
- Urodynamics Teaching Consult LEB
Topic: Clean Intermittent Catheterization





Physician Orders PEDIATRIC: LEB UROL General Admit Plan

Consult Wound Care Nurse

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

***Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

