



Physician Orders ADULT: LEB Fecal Microbiota Transplant Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- Initiate Powerplan Phase
T;N, Phase: LEB Fecal Microbiota Transplant Phase, When to Initiate:

LEB Fecal Microbiota Transplant Phase

Admission/Transfer/Discharge

- Patient Status Initial Outpatient
T;N Attending Physician:
Reason for Visit:
Bed Type: Specific Unit:
Outpatient Status/Service: OP-Ambulatory Surgery
Notify Physician-Once
T;N, of room number on arrival to unit.

Food/Nutrition

- NPO
Start at: T;23:59, after midnight (DEF)\*
Start at: T;N

Patient Care

- Consent Signed For
T;N, Procedure: Fecal Microbiota Transplant, Utilize specific Consent for Fecal Microbiota Transplantation (FMT) located under Clinical and System Policies on MOLLI.
NGT Insert
T;N, for Fecal Microbiota Transplant procedure if not already in place. Placement must be confirmed by x-ray.
IV Insert/Site Care
T;N, for Fecal Microbiota Transplant procedure if not already in place.

Nursing Communication

- Nursing Communication
T;N, Discontinue antibiotics at midnight the night before the procedure.
Nursing Communication
T;N, If NGT is placed, enter order for KUB stat to confirm placement before procedure

Continuous Infusion

- Sodium Chloride 0.9%
1,000 mL, IV, Routine, (for 24 hr ), TKO
Comments: To Keep Vein Open: Do not infuse more than one bag or 1000 mL.
Dextrose 5% in Water
1,000 mL, IV, Routine, (for 24 hr ), TKO





**Physician Orders ADULT: LEB Fecal Microbiota Transplant Plan**

*Comments: To Keep Vein Open: Do not infuse more than one bag or 1000 mL.*

- Sodium Chloride 0.45%  
1,000 mL, IV, Routine, (for 24 hr ), TKO

*Comments: To Keep Vein Open: Do not infuse more than one bag or 1000 mL.*

**Medications**

If Fecal Microbiota Transplant is delivered via lower route, order drug below to be given post procedure(NOTE)\*

- loperamide  
0.5 mg/kg, Liq, PO, bid, Routine, (for 2 dose )  
*Comments: To be given post-procedure*

- polyethylene glycol 3350 with electrolytes  
25 mL/kg, Oral Soln, PO, q1h, Routine, (for 4 dose )  
*Comments: To be given for 4 doses or until rectal effluent is clear. To be given the night before procedure.*

If Fecal Microbiota Transplant is delivered via nasogastric tube, order pantoprazole below.(NOTE)\*

- pantoprazole  
1 mg/kg, Injection, IV Push, N/A, Routine  
*Comments: Administer the evening before the procedure at 20:00. Max dose: 40 mg*

- pantoprazole  
1 mg/kg, Injection, IV Push, N/A, Routine  
*Comments: Administer the morning of the procedure at 06:30. Max dose: 40 mg*

**Laboratory**

- Electrolytes  
*Routine, T;N, once, Type: Blood*
- Fecal Microbiota 250 mL Instillation Product  
*Routine, FMT Delivery Mode Colonoscopy*
- Fecal Microbiota 250 mL Instillation Product  
*Routine, FMT Delivery Mode Lower Delivery Enema*
- Fecal Microbiota 30 mL Instillation Product  
*Routine, FMT Delivery Mode NGT Tube*
- Fecal Microbiota 30 mL Instillation Product  
*Routine, FMT Delivery Mode Nasal Duodenal Tube*
- Fecal Microbiota 30 mL Instillation Product  
*Routine, FMT Delivery Mode Upper Delivery EGD*
- Instill Fecal Microbiota
  - Routine, FMT Delivery Mode Colonoscopy (DEF)\**
  - Routine, FMT Delivery Mode Lower Delivery Enema*





Physician Orders ADULT: LEB Fecal Microbiota Transplant Plan

- Routine, FMT Delivery Mode NGT Tube
- Routine, FMT Delivery Mode Nasal Duodenal Tube
- Routine, FMT Delivery Mode Upper Delivery EGD

Consults/Notifications/Referrals

- GI Lab Request To Schedule
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Colonoscopy with anesthesia. (DEF)\*
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Colonoscopy without anesthesia
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Colonoscopy with anesthesia
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Colonoscopy without anesthesia
- GI Lab Request To Schedule
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Enema (DEF)\*
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For lower delivery Enema
- GI Lab Request To Schedule
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery EGD with anesthesia (DEF)\*
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery EGD without anesthesia
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery EGD with anesthesia
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery EGD without anesthesia
- GI Lab Request To Schedule
  - T+1, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery via NGT (DEF)\*
  - T;N, Procedure: Fecal Microbiota Transplant (FMT) For upper delivery via NGT
- Infection Control Consult
  - T;N, Reason for Consult: Fecal Microbiota Transplant

|      |      |                       |           |
|------|------|-----------------------|-----------|
| Date | Time | Physician's Signature | MD Number |
|------|------|-----------------------|-----------|

**\*Report Legend:**  
 DEF - This order sentence is the default for the selected order  
 GOAL - This component is a goal





**Physician Orders ADULT: LEB Fecal Microbiota Transplant Plan**

IND - This component is an indicator  
INT - This component is an intervention  
IVS - This component is an IV Set  
NOTE - This component is a note  
Rx - This component is a prescription  
SUB - This component is a sub phase, see separated sheet  
R-Required order

