



Physician Orders PEDIATRIC: LEB ED Vomiting/Diarrhea Plan

LEB ED Standing Orders Vomiting/Diarrhea

Non Categorized

Criteria: Patients < 18 years of age with vomiting/diarrhea of recent onset not due to chronic disease with or without fever, nausea, or abdominal pain.(NOTE)*

Vital Signs

- Vital Signs
T;N, Stat Monitor and Record T,P,R,BP, per routine

Food/Nutrition

- NPO
Start at: T;N

Patient Care

- IV Insert/Site Care LEB
T;N, Stat, q2h(std)

Laboratory

- CBC
STAT, T;N, once, Type: Blood
- BMP
STAT, T;N, once, Type: Blood
- If possibility of pregnancy, place order below:(NOTE)*
- Pregnancy Screen Serum
STAT, T;N, once, Type: Blood

LEB ED Vomiting/ Diarrhea Plan Phase

Non Categorized

- Powerplan Open

Patient Care

- Bedside Glucose Nsg
T;N, Stat
- PO Challenge
T;N, Stat
- O2 Sat Spot Check-NSG
T;N, Stat, with vitals
- O2 Sat Monitoring NSG
T;N, Stat
- Cardiopulmonary Monitor
T;N Stat, Monitor Type: CP Monitor

Respiratory Care

- ISTAT POC (RT Collect)
*T;N Stat once, Test Select Arterial Blood Gas (DEF)**
T;N Stat once, Test Select Venous Blood Gas





Physician Orders PEDIATRIC: LEB ED Vomiting/Diarrhea Plan

- Oxygen Delivery
T;N, Special Instructions: Titrate to keep O2 sat \geq 92%. Wean to room air.

Continuous Infusion

- Sodium Chloride 0.9% Bolus
 - 20 mL/kg, IV, once, STAT, (infuse over 15 min), (Bolus) (DEF)*
 - 10 mL/kg, IV, once, STAT, (infuse over 15 min), (Bolus)
 - 10 mL/kg, IV, once, STAT, (infuse over 30 min), (Bolus)
 - 20 mL/kg, IV, once, STAT, (infuse over 30 min), (Bolus)
- Sodium Chloride 0.9%
1,000 mL, IV, STAT, mL/hr
- D5 1/2NS
1,000 mL, IV, STAT, mL/hr
- D5 1/4 NS
1,000 mL, IV, STAT, mL/hr
- D5 1/2 NS KCl 20 mEq/L
1,000 mL, IV, STAT, mL/hr
- D5 1/4 NS KCl 20 mEq/L
1,000 mL, IV, STAT, mL/hr

Medications

- metoclopramide
0.1 mg/kg, Ped Injectable, IV Push, once, STAT, Max: 10 mg/dose
- ondansetron
 - 0.1 mg/kg, Oral Susp, PO, once, STAT, Max dose = 4 mg (DEF)*
 - 4 mg, Orally Disintegrating Tab, PO, once, STAT
- ondansetron
0.1 mg/kg, Ped Injectable, IV Push, once, STAT, Max dose = 8 mg

Laboratory

- Urinalysis w/Reflex Microscopic Exam
STAT, T;N, Type: Urine, Nurse Collect
- Stool Culture
STAT, T;N, Specimen Source: Stool, Nurse Collect
- WBC Stool - Ped
STAT, T;N, once, Type: Stool, Nurse Collect
- Stool Ova & Parasites
STAT, T;N, Specimen Source: Stool Rectum, Nurse Collect, Method: Stool
- Rotavirus Antigen
STAT, T;N, once, Type: Stool, Nurse Collect





Physician Orders PEDIATRIC: LEB ED Vomiting/Diarrhea Plan

- Pregnancy Screen Serum
STAT, T;N, once, Type: Blood

Diagnostic Tests

- Abd Flat and Upright/Decubitus
T;N, Reason for Exam: Nausea and Vomiting, Stat, Stretcher

Date

Time

Physician's Signature

MD Number

***Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

