



## Physician Orders ADULT: Head and Neck Postoperative Plan

### Initiate Orders Phase

#### Care Sets/Protocols/PowerPlans

- ☐ Initiate Powerplan Phase

*T;N, Phase: Head and Neck Postoperative Phase, When to Initiate: \_\_\_\_\_*

### Head and Neck Post Operative Phase

#### Vital Signs

- ☒ Vital Signs

*T;N, Monitor and Record T,P,R,BP, per postop routine special instructions: P, R, BP q 30 minutes x2, then q 1 hour x2, then q4hr x24 hours. Temp q4hr.*

- ☐ Vital Signs

☐ *T;N, Monitor and Record T,P,R,BP, q2h(std) (DEF)\**

☐ *T;N, Monitor and Record T,P,R,BP, q4h(std)*

☐ *T;N, Monitor and Record T,P,R,BP, q8h(std)*

#### Activity

- ☐ Bedrest

*T;N, Routine*

- ☐ Activity As Tolerated

*T;N*

- ☐ Up To Chair

*T;N*

#### Food/Nutrition

- ☐ NPO

*Start at: T;N*

- ☐ Clear Liquid Diet

*Start at: T;N, Adult (>18 years)*

- ☐ Full Liquid Diet

*Start at: T;N, Adult (>18 years)*

- ☐ GI Soft Diet

*Start at: T;N*

- ☐ Regular Diet

*Start at: T;N*

- ☐ AHA Diet

*Start at: T;N, Adult (>18 years)*

- ☐ Renal Diet Not On Dialysis

*Start at: T;N, Adult (>18 years)*

- ☐ Renal Diet On Dialysis

*Start at: T;N*

- ☐ Consistent Carbohydrate Diet

*Start at: T;N, 1800 Calorie*

- ☐ Tube Feeding Continuous/Int Plan(SUB)\*

- ☐ Tube Feeding Titrate Plan(SUB)\*





**Physician Orders ADULT: Head and Neck Postoperative Plan**

- ☐ Tube Feeding Bolus Plan(SUB)\*

**Patient Care**

- ☐ Drain Care  
*T;N, Jackson-Pratt(s) to bulb suction*
- ☐ Drain Care  
*T;N, q-shift, charge, PJP Drains-empty and recharge and record output*
- ☐ Elevate Head Of Bed  
*T;N, 30 degrees*
- ☐ Foley Care  
*T;N*
- ☒ Incentive Spirometry NSG  
*T;N, q1h-Awake*
- ☒ O2 Sat Continuous Monitoring NSG  
*T;N*
- ☐ Trach Care  
*T;N, Change inner cannula q 12 hrs/prn, if obstructed*
- ☐ Trach Care  
*T;N, Keep same size Trach and smaller size Trach at bedside*
- ☐ Trach Care  
*T;N, Do not change trach ties unless okay with MD*
- ☐ Trach Care  
*T;N, Place obturator at head of bed*
- ☐ Trach Care  
*T;N, q8h(std)*
- ☐ Suction Patient  
*T;N, q2h(std), Suctioning Type: Trach*
- ☐ Suction Patient  
*T;N, prn, PRN*
- ☐ Nursing Communication  
*T;N, q12h, clean the stoma under trach flange with Q-tips/gauze soaked in NS*
- ☐ Mouth Care  
*T;N, Routine, Clean oral cavity with antiseptic oral care kit*
- ☐ Accucheck Nsg  
*T;N, q6h(std)*

**Respiratory Care**

- ☐ O2-BNC  
*T;N, 2 L/min*
- ☐ O2-Trach Collar  
*T;N*

**Continuous Infusion**

- ☒ D5 1/2 NS KCl 20 mEq/L





**Physician Orders ADULT: Head and Neck Postoperative Plan**

- ☐ 1,000 mL, IV, 100 mL/hr
- ☐ D5 1/2NS
- ☐ 1,000 mL, IV, 100 mL/hr
- ☐ Sodium Chloride 0.9%
- ☐ 1,000 mL, IV, 100 mL/hr
- ☐ Sodium Chloride 0.45%
- ☐ 1,000 mL, IV, 100 mL/hr

**Medications**

- ☒ VTE SURGICAL Prophylaxis Plan(SUB)\*
- ☐ PCA - MorPHINE Protocol Plan (Adult)(SUB)\*
- ☐ PCA - HYDROmorphine Protocol Plan (Adult)(SUB)\*
- ☐ Insulin SENSITIVE Sliding Scale Plan(SUB)\*
- ☐ Insulin STANDARD Sliding Scale Plan(SUB)\*
- ☐ Insulin RESISTANT Sliding Scale Plan(SUB)\*
- Please enter the Start Time on the antibiotic orders below based on time of preop dose(NOTE)\*
- If antibiotics are being continued for greater than 24 hrs, please document your reason(NOTE)\*
- ☐ **+1 Hours** ceFAZolin
- 2 g, Injection, IV Piggyback, once
- Comments: Time dose 8 hours after pre op dose
- If patient has penicillin or cephalosporin allergy place BOTH orders below(NOTE)\*
- ☐ **+1 Hours** clindamycin
- 600 mg, IV Piggyback, IV Piggyback, once
- Comments: Time dose 8 hours after pre op dose
- AND(NOTE)\*
- ☐ **+1 Hours** gentamicin
- 2 mg/kg, Injection, IV Piggyback, once
- Comments: Time dose 12 hours after the pre op dose
- ☒ **+1 Hours** ondansetron
- 4 mg, Injection, IV Piggyback, q6h, PRN Nausea/Vomiting
- ☒ **+1 Hours** bacitracin 500 units/g topical ointment
- 1 application, Ointment, TOP, bid
- Choose one of the below pain medications for Mild-Moderate pain(NOTE)\*
- ☐ **+1 Hours** acetaminophen-oxyCODONE 325 mg-5 mg oral tablet
- 1 tab, Tab, PO, q4h, PRN Pain, Moderate (4-7)
- ☐ **+1 Hours** acetaminophen-HYDROcodone 325 mg-7.5 mg/15 mL oral solution
- 0.15 mg/kg, Elixir, PO, q4h, PRN Pain, Moderate (4-7)
- Choose one of the below pain medications for SEVERE pain(NOTE)\*
- ☐ **+1 Hours** morphine
- 1 mg, Injection, IV Push, q2h, PRN Pain, Severe (8-10)
- Comments: hold if RR less than 12
- ☐ **+1 Hours** HYDROmorphine
- 0.5 mg, Injection, IV Push, q2h, PRN Pain, Severe (8-10)





## Physician Orders ADULT: Head and Neck Postoperative Plan

*Comments: hold if RR less than 12*

### Laboratory

- ☐ CBC  
*Routine, T;N, once, Type: Blood*
- ☐ BMP  
*Routine, T;N, once, Type: Blood*
- ☐ Magnesium Level  
*Routine, T;N, once, Type: Blood*
- ☐ Phosphorus Level  
*Routine, T;N, once, Type: Blood*
- ☐ Ionized Calcium  
*Routine, T;N, once, Type: Blood*
- ☐ CBC  
*Routine, T+1;0400, once, Type: Blood*
- ☐ BMP  
*Routine, T+1;0400, once, Type: Blood*
- ☐ Magnesium Level  
*Routine, T+1;0400, once, Type: Blood*
- ☐ Phosphorus Level  
*Routine, T+1;0400, once, Type: Blood*
- ☐ Ionized Calcium  
*Routine, T+1;0400, once, Type: Blood*
- ☐ PTH Profile  
*Routine, T;N+240, once, Type: Blood*  
*Comments: 4 hours Post op*

### Diagnostic Tests

- ☐ Chest 1VW Frontal  
*T;N, Reason for Exam: Other, Enter in Comments, Stat, Portable*  
*Comments: Reason: Post trach(PACU ORDER)*
- ☐ Chest 1VW Frontal  
*T+1;0800, Reason for Exam: Other, Enter in Comments, Routine*  
*Comments: Reason: Status Post trach*
- ☐ KUB  
*T;N, Reason for Exam: Other, Enter in Comments, Stat, Portable*  
*Comments: Reason: NG Tube placement(PACU Order)*

### Consults/Notifications/Referrals

- ☒ Notify Physician For Vital Signs Of  
*T;N, BP Systolic > 180, BP Diastolic > 100, BP Systolic < 100, Celsius Temp > 38.5, Oxygen Sat < 92, Urine Output < 30*
- ☐ Dietitian Consult/Nutrition Therapy  
*T;N, Type of Consult: Enteral/Tube Feeding*
- ☐ Speech Therapy Initial Eval and Tx





**Physician Orders ADULT: Head and Neck Postoperative Plan**

*T;N, Routine*

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

**\*Report Legend:**

DEF - This order sentence is the default for the selected order  
GOAL - This component is a goal  
IND - This component is an indicator  
INT - This component is an intervention  
IVS - This component is an IV Set  
NOTE - This component is a note  
Rx - This component is a prescription  
SUB - This component is a sub phase, see separate sheet  
R-Required order

