



**Physician Orders ADULT: VTE ORTHO SURGICAL Prophylaxis Plan**

**VTE ORTHO SURGICAL Prophylaxis Plan**

**Non Categorized**

Bleeding Risk Factor Assessment criteria is listed below VTE orders.(NOTE)\*

**Do Not Administer VTE Prophylaxis**

If both Mechanical and Pharmacological VTE prophylaxis is contraindicated, place order below:(NOTE)\*

- Reason Surgical VTE Prophylaxis Not Received

**Ortho Procedures**

NOTE: If bleeding risk exists and NO contraindications to SCDs, place order below:(NOTE)\*

- Sequential Compression Device Apply  
*T;N, Apply To Lower Extremities, Bleeding Risks Present*

If No bleeding risk exists, order one of the medications AND BOTH CBC w/o Diff orders below.(NOTE)\*

- +720 Minutes** fondaparinux  
*2.5 mg, Injection, Subcutaneous, QDay, Routine, Do not adjust time of first dose as scheduled by pharmacy.*

OR(NOTE)\*

- +720 Minutes** enoxaparin  
*30 mg, Injection, Subcutaneous, bid, Routine*  
*Comments: First dose should be timed to be given at least 12 hours post op, If CrCl less than 30mL/min, pharmacy to adjust dose to 30mg SQ QDay. Do not adjust time of first dose as scheduled by pharmacy.*

OR(NOTE)\*

- +480 Minutes** rivaroxaban  
*10 mg, Tab, PO, q24h, Routine*  
*Comments: first dose should be timed to be given at least 6-10 hours post op, If CrCl less than 30mL/min, use is contraindicated. Pharmacist may adjust administration times after first dose. If the patient had an AM surgery, the first dose should be given that day at 2200. If the patient had a PM surgery, the first dose should be given at 0600 the following day.*

- +720 Minutes** apixaban  
*2.5 mg, Tab, PO, bid, Routine*  
*Comments: first dose should be timed to be given at least 12 hours post op, If CrCl less than 25mL/min, use is contraindicated. If the patient had an AM surgery, the first dose should be given that day at 2200. If the patient had a PM surgery, the first dose should be given at 0600 the following day.*

OR(NOTE)\*

Aspirin is an option for VTE Prophylaxis for patients with hip or knee surgeries only.(NOTE)\*

- +240 Minutes** aspirin
  - 81 mg, DR Tablet, PO, QDay, First dose should be timed to be given at least 4-6 hours post op. (DEF)\**  
*Comments: "Aspirin For VTE Prophylaxis".*
  - 81 mg, DR Tablet, PO, bid, First dose should be timed to be given at least 4-6 hours post op. Comments: "Aspirin For VTE Prophylaxis".*







**Physician Orders ADULT: VTE ORTHO SURGICAL Prophylaxis Plan**

**\*Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

