

Physician Orders ADULT

Order Set: FLAG-IDA

Diagnosis : AML				
Height		kg		Cycle: Of :
Actual			m2	Day/Wk: Freq:
Allergies: [] No known allergies				
[]Medication allergy(s):				
[] Latex allergy []Other: Patient Care				
[]	Nursing Communication T;N, Do not exceed a treatment BSA of m2			
[]	Nursing Communication	T;N, May hold hydration during chemotherapy infusion		
[]	Nursing Communication	T;N, Verify patient has had MUGA or ECHO to r/o Cardiac dysfunction prior to chemotherapy		
Continuous Infusions				
Pre Hydration				
	Normal Saline 1,000 mL, IV, Routine,mL/hr			
Medications				
[X]	prednisoLONE opthalmic 1%	2 drops, Opthalmic Susp, Both eyes, q6h, on DAYS 1-7		
[X]	filgrastim	5 mcg/kg, Injection, SQ, daily, Start on DAY 6 Comment : Give until ANC greater than 1000 for 2 days		
CHEMOTHERAPY				
	Drug (generic) & solution (optional)	Intended Dose	Actual Dose	Route, Infusion, Frequency and total doses
[X]	fludarabine	30 mg/m ²		IV Piggyback, Infuse over 30 min, q24h on DAYS 1-5
[X]	cytarabine	2000 mg/m ²		IV Piggyback, Infuse over 3 hours, q24h on DAYS 1-5 , give 4 hours after the fludarabine
[X]	IDArubicin	10 mg/m ²		IV Push, Daily on DAYS 1- 3
Acute Emesis Prophylaxis (may undergo therapeutic interchange)				
NOTE: Administer intial doses at least 30-60 minutes prior to chemotherapy				
[X]	ondansetron	12 mg, Injection, IV Piggyback, qDay, on DAYS 1-5		
[X]	dexamethasone	8 mg, Injection, IV Push, Q Day , on DAYS 1 - 5		
[X]	prochlorperazine	10 mg, Tab, PO, q6h, PRN Nausea/Vomiting		
[X]	prochlorperazine	10 mg, Injection, IV Push, q6h, PRN Nausea/Vomiting , Comment : if unable to take PO		
Consults/Notifications				
[]	Notify Physician- Once T;N, Who:, For: if BSA exceeds 2 m ²			

Time

Physician's Signature

MD Number