



Physician Orders ADULT: Gastroenteritis Observation Plan

Initiate Orders Phase

Admission/Transfer/Discharge

- ☒ Patient Status Initial Outpatient
T;N Attending Physician: _____
Reason for Visit: _____
Bed Type: _____ Specific Unit: _____
Outpatient Status/Service: Outpatient Status/Service OP-OBSERVATION Services

Care Sets/Protocols/PowerPlans

- ☒ Initiate Powerplan Phase
Phase: Gastroenteritis Observation Phase, When to Initiate: _____

Gastroenteritis Observation Phase

Vital Signs

- ☐ Vital Signs
Routine Monitor and Record T,P,R,BP, q4h(std)
- ☐ Orthostatic Blood Pressure
Stat, with heart rate

Activity

- ☐ Bedrest
Routine
- ☐ Bedrest w/BRP
- ☐ Out Of Bed

Food/Nutrition

- ☐ NPO
- ☐ Clear Liquid Diet
Adult (>18 years)
- ☐ Full Liquid Diet
- ☐ Regular Adult Diet
- ☐ Consistent Carbohydrate Diet
Caloric Level: 1800 Calorie
- ☐ Gastroenteritis Diet

Patient Care

- ☒ Intermittent Needle Therapy Insert/Site Care
Stat, q4day
- ☒ Instruct/Educate
Instruct: Patient and Family, Method: Provide Pamphlet, Topic: Observation Services
- ☒ Intake and Output
- ☐ Weight
once, Obtain Actual Weight
- ☐ Nasogastric Tube
Insert and put to low continuous suction
- ☐ O2 Sat Continuous Monitoring NSG
Routine

Continuous Infusion

- ☐ Sodium Chloride 0.9% Bolus
1,000 mL, IV Piggyback, IV Piggyback, once, STAT, 1,000 mL/hr
- ☐ Lactated Ringers Injection
1,000 mL, IV, Routine, 75 mL/hr
- ☐ Dextrose 5% in Lactated Ringers Injection
1,000 mL, IV, Routine, 75 mL/hr
- ☐ Dextrose 5% with 0.45% NaCl





Physician Orders ADULT: Gastroenteritis Observation Plan

- ☐ 1,000 mL, IV, Routine, 75 mL/hr
- ☐ Sodium Chloride 0.9%
1,000 mL, IV, Routine, 75 mL/hr
- ☐ Dextrose 5% NaCl 0.45% KCl 20 mEq
20 mEq 1,000 mL, IV, Routine, 75 mL/hr

Medications

- ☐ **+1 Hours** metoclopramide
10 mg, Injection, IV Push, q6h, Nausea/Vomiting
- ☐ **+1 Hours** acetaminophen
650 mg, Tab, PO, q6h, PRN Pain, Mild (1-3), Routine
- ☐ **+1 Hours** acetaminophen
650 mg, Supp, PR, q4h, PRN Pain, Mild or Fever, Routine
- ☐ **+1 Hours** acetaminophen-HYDROcodone 325 mg-5 mg oral tablet
1 tab, Tab, PO, q6h, PRN Pain, Moderate (4-7), Routine
- ☐ **+1 Hours** promethazine
12.5 mg, Tab, PO, q4h, PRN Nausea, Routine
- ☐ **+1 Hours** ondansetron
4 mg, Injection, IV Push, q8h, PRN Nausea/Vomiting, Routine
- ☐ **+1 Hours** temazepam
7.5 mg, Cap, PO, hs, PRN Sleep, Routine
Comments: May repeat dose once after 30 min if required.
- ☐ **+1 Hours** pantoprazole
 - ☐ 40 mg, Cap, PO, QDay, Routine (DEF)*
 - ☐ 40 mg, Injection, IV Push, q24h, Routine
- ☐ **+1 Hours** famotidine
20 mg, Injection, IV Push, q12h, Routine, Change to q24h if CrCl is less than 50mL/min
Comments: Change to q24h if CrCl is less than 50 mL/min

Laboratory

- ☐ CBC
STAT, T;N, once, Type: Blood
- ☐ CMP
STAT, T;N, once, Type: Blood
- ☐ BMP
STAT, T;N, once, Type: Blood
- ☐ Calcium Ionized
STAT, T;N, once, Type: Blood
- ☐ Magnesium Level
STAT, T;N, once, Type: Blood
- ☐ Phosphorus Level
STAT, T;N, once, Type: Blood
- ☐ Amylase Level
STAT, T;N, once, Type: Blood
- ☐ Lipase Level
STAT, T;N, once, Type: Blood
- ☐ Hepatitis Profile (A,B & C)
Routine, T;N, once, Type: Blood
- ☐ Hepatic Panel
Routine, T;N, once, Type: Blood
- ☐ Urinalysis w/Reflex Microscopic Exam
STAT, T;N, once, Type: Urine, Nurse Collect





Physician Orders ADULT: Gastroenteritis Observation Plan

- ☐ Clostridium difficile Toxin B gene by PCR
Routine, T;N, once, Type: Stool, Nurse Collect
- ☐ Ova & Parasites Stool
Routine, T;N, Specimen Source: Stool, Nurse Collect
- ☐ Stool Culture
Routine, T;N, Specimen Source: Stool, Nurse Collect

Diagnostic Tests

- ☐ Chest 1 View
T;N, Routine, Portable
- ☐ Chest 2 Views
T;N, Routine, Stretcher
- ☐ Abdomen 2 Views
T;N, Routine, Stretcher
- ☐ CT Abdomen & Pelvis W/Cont Plan(SUB)*

Consults/Notifications/Referrals

- ☒ Notify Physician-Once
Notify For: room number on arrival to unit
- ☐ Physician Consult
- ☐ Physician Consult
- ☐ Case Management Consult
Routine, Reason: Discharge Planning

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

*Report Legend:

DEF - This order sentence is the default for the selected order
 GOAL - This component is a goal
 IND - This component is an indicator
 INT - This component is an intervention
 IVS - This component is an IV Set
 NOTE - This component is a note
 Rx - This component is a prescription
 SUB - This component is a sub phase, see separate sheet
 R-Required order

