



Physician Orders PEDIATRIC: LEB Croup Admit Plan

Initiate Orders Phase

Care Sets/Protocols/PowerPlans

- Initiate Powerplan Phase
Phase: LEB Croup Admit Phase, When to Initiate: _____

LEB Croup Admit Phase

Admission/Transfer/Discharge

- Patient Status Initial Outpatient
*T;N Attending Physician: _____
Reason for Visit: _____
Bed Type: _____ Specific Unit: _____
Outpatient Status/Service: [] Ambulatory Surgery, [] OP Diagnostic Procedure
[] OP OBSERVATION Services*
- Notify Physician-Once
Notify For: of room number on arrival to unit

Vital Signs

- Vital Signs
 - Monitor and Record T,P,R,BP, q4h(std) (DEF)*
 - Monitor and Record T,P,R,BP

Activity

- Activity As Tolerated
Up Ad Lib

Food/Nutrition

- NPO
- Breastfeed
- LEB Formula Orders Plan(SUB)*
- Regular Pediatric Diet
- Lactating Mother Guest Tray

Patient Care

- Advance Diet As Tolerated
start clear liquids and advance to regular diet as tolerated
- Isolation Precautions
Isolation Type: Droplet Precautions (DEF)
per policy*
- Intake and Output
Routine, q2h(std)
- Daily Weights
Routine, qam





Physician Orders PEDIATRIC: LEB Croup Admit Plan

- Suction Patient
q2h(std), PRN, Suction: Nasal, nasopharyngeal suction only when necessary
- O2 Sat Spot Check-NSG
with vital signs
- O2 Sat Monitoring NSG
- Cardiopulmonary Monitor
Routine, Monitor Type: CP Monitor

Respiratory Care

- Oxygen Delivery
Special Instructions: Titrate to keep O2 sat \geq 92%. Wean to room air.

Continuous Infusion

- Sodium Chloride 0.9%
1,000 mL, IV, Routine, mL/hr
- D5 1/2NS
1,000 mL, IV, Routine, mL/hr
- D5 1/2 NS KCl 20 mEq/L
1,000 mL, IV, Routine, mL/hr

Medications

- +1 Hours** racepinephrine 2.25% inhalation solution
0.25 mL, Inh Soln, NEB, q4h, PRN Stridor, Routine
- dexamethasone
 - 0.6 mg/kg, Ped Injectable, IV Piggyback, once, Routine, [Less than 25 kg], For Croup (DEF)**
 - 16 mg/kg, Ped Injectable, IV Piggyback, once, Routine, [Greater than 25 kg], For Croup*
- +1 Hours** dexamethasone
 - 4 mg, Tab, PO, once, Routine, [7-10.9 kg], For Croup (DEF)**
 - 8 mg, Tab, PO, once, Routine, [11-14.9 kg], For Croup*
 - 12 mg, Tab, PO, once, Routine, [15-24.9 kg], For Croup*
 - 16 mg, Tab, PO, once, Routine, [Greater than 25 kg], For Croup*
- +1 Hours** acetaminophen
 - 10 mg/kg, Liq, PO, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day (DEF)**
 - 80 mg, Chew tab, PO, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day*
- +1 Hours** acetaminophen
10 mg/kg, Supp, PR, q4h, PRN Pain, Mild or Fever, Routine, Max Dose = 75 mg/kg/day up to 4g/day
Comments: May give PR if unable to take PO





Physician Orders PEDIATRIC: LEB Croup Admit Plan

Laboratory

- CBC
T;N, Routine, once, Type: Blood
- CMP
Routine, T;N, once, Type: Blood
- BMP
Routine, T;N, once, Type: Blood
- RSV Antigen Screen
Routine, T;N, once, Type: Nasopharyngeal(N-P), Nurse Collect
- Respiratory Virus Panel by PCR LEB
Routine, T;N, once, Type: Nasopharyngeal(N-P), Nurse Collect
- Influenza A/B Antigen
Routine, T;N, once, Type: Nasopharyngeal(N-P)
- Influenza/RSV Panel PCR
Routine, T;N, once, Type: Nasopharyngeal(N-P), Nurse Collect
- Bordetella pertussis by PCR
Routine, T;N, once, Type: Nasopharyngeal(N-P), Nurse Collect

Diagnostic Tests

- Neck Soft Tissue
T;N, Reason for Exam: Other, Enter in Comments, ROUTINE, Wheelchair
Comments: croupy cough

Consults/Notifications/Referrals

- Consult MD Group
- Consult MD
- Notify Physician-Continuing
Notify For: Respiratory distress, increased oxygen requirements

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

*Report Legend:

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention



Attach patient label here



Physician Orders PEDIATRIC: LEB Croup Admit Plan

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase, see separate sheet

R-Required order

