

Height: \_\_\_\_\_ cm Weight: \_\_\_\_\_ kg

## Palivizumab (Synagis) Order Form

Please allow 48 hours for verification and dispensing.

1. Palivizumab (15 mg/kg) mg IM x one dose.

Patients must meet at least one of the following criteria, never have been home, and have plans for discharge within 1 week. \*

2. Please select indication for use:

 $\leq$  32 weeks EGA infant who is less than 6 months of age

 $\leq$  29 weeks EGA infant who is less than 1 year of age

Between 32-35 weeks EGA infant who is less than 6 months of age in October with additional risk factors. Additional Risk factors include:

day care attendance

multiple <8 year old children in household

tobacco smoke in home

severe neuromuscular disease

congenital malformations affecting the respiratory tract

Infant with Chronic Lung Disease who is less than 2 years of age in October if: systemic steroid, oxygen, or diuretic dependent within the last 6 months.

Infant with <u>severe</u> immunodeficiency (ex: recent transplantation, SCID, prolonged high-intensity chemotherapy) who is less than 2 years of age in October if significant exposure to RSV is likely.

Infant less than 2 years of age with hemodynamically significant congenital heart disease as judged by the patient's pediatric cardiologist.

\* Patients who have previously been home should receive their palivizumab from their PCP. If necessary, the PCP can send the patient's dose to the hospital to be dispensed by the pharmacy. The Infectious Diseases Service should be contacted for use outside the above criteria.

Physician Name		Date	Time
----------------	--	------	------

Physician Signature Physician #