



Physician Orders ADULT ED Pre Eclampsia/Eclampsia Orders

[R] = will be ordered

T= Today; N = Now (date and time ordered)

Height: _____ cm Weight: _____ kg

Allergies:		
<input type="checkbox"/> No known allergies		
<input type="checkbox"/> Medication allergy(s): _____		
<input type="checkbox"/> Latex allergy <input type="checkbox"/> Other: _____		
Vital Signs		
<input type="checkbox"/>	Vital Signs	T;N, Include: BP, HR, RR, DTRs, O2 Sat, LOC, breath sounds
<input type="checkbox"/>	Fetal Heart Tones Auscultation (FHR Assessment)	Routine, Continuous fetal heart rate and uterine contraction monitoring per Magnesium Sulfate Administration Policy.
Activity		
<input type="checkbox"/>	Bedrest	T;N
Food/Nutrition		
<input type="checkbox"/>	NPO	Start at: T;N
<input type="checkbox"/>	NPO	T;N, except for ice chips and medications
Patient Care		
<input type="checkbox"/>	Intake and Output	q1h
<input type="checkbox"/>	Indwelling Urinary Catheter Insert-Follow Removal Protocol	bag with urometer
<input type="checkbox"/>	Indwelling Urinary Catheter Care	
<input type="checkbox"/>	Seizure Precautions	
<input type="checkbox"/>	O2 Sat Continuous Monitoring NSG	
<input type="checkbox"/>	Nursing Communication	Notify physician immediately and Stop magnesium sulfate infusion for symptoms of magnesium toxicity including double or blurred vision, tachycardia or bradycardia, respiratory rate < 14 or > 24/min, oxygen saturation < 95%, changes in breath sounds (r/o pulmonary edema), changes in level of consciousness, or loss of deep tendon reflexes.
<input type="checkbox"/>	Nursing Communication	T;N, Upon completion of magnesium sulfate bolus, place order for magnesium level q6h while receiving magnesium.
Continuous Infusions		
<input type="checkbox"/>	Lactated Ringers	1,000 mL, IV, Routine, T;N, 50 mL/hr, Comments: titrate total IV fluid volume to total 100 mL/hr
<input type="checkbox"/>	magnesium sulfate 20 g/ LR infusion	20 g / 500 mL, IV, Routine, T;N, 50 mL/hr, T;N, Comment: Initial Rate: 50mL/hr =2g/hr, MAINTENANCE dose
Medications		
<input checked="" type="checkbox"/>	magnesium sulfate	6 g, Injection, IV Piggyback, once, Routine, T;N, (infuse over 30 min), (OB patient ONLY), LOADING dose, Comment: Give via infusion pump in hub nearest to patient
<input type="checkbox"/>	magnesium sulfate	4 g, Injection, IV Piggyback, once, Routine, T;N, (infuse over 30 min), (OB patient ONLY), LOADING dose, Comment: Infuse via infusion pump in hub nearest patient.
Medications		
<input type="checkbox"/>	magnesium sulfate	6 g, Injection, IV Piggyback, N/A (1 dose), PRN seizure activity, Routine, T;N, (infuse over 30 min), (OB patient ONLY), Comment: Infuse via infusion pump in hub nearest to patient
<input type="checkbox"/>	LORazepam	2 mg, Injection, IV Push, N/A (1 dose), PRN seizure activity, Routine, T;N, Comment: for persistent seizure activity not resolved by PRN magnesium bolus
<input type="checkbox"/>	labetalol	20 mg, Injection, IV Push, q15min X 2, PRN Hypertension, Comment :Give for 2 or more instances of Systolic >160 and/ or Diastolic >110
<input type="checkbox"/>	calcium gluconate	1g, Injection, IV Push, N/A (1 dose), PRN signs & symptoms of magnesium toxicity, Routine, T;N, Comment: Administer with MD Supervision



