



Physician Orders PEDIATRIC: LEB ED Non-Critical Trauma Plan

LEB ED Triage Orders Non Critical Trauma

Non Categorized

Criteria: Patients between 0-18 years of age who present with trauma and do not meet the LeBonheur Trauma Stat or Trauma Alert Guidelines.(NOTE)*

Vital Signs

- Vital Signs
Monitor and Record T,P,R,BP, Per ED policy

Food/Nutrition

- NPO

Patient Care

- O2 Sat Monitoring NSG
q2h(std)
- Cardiopulmonary Monitor
Stat, Monitor Type: O2 Monitor
- Nursing Communication
If there is a suspected fracture, contact the physician to obtain a diagnostic order.
- Nursing Communication
Apply moist saline bandages to any deep abrasion or laceration

LEB ED Non-Critical Trauma Phase

Non Categorized

R Powerplan Open

Patient Care

- LEB ED Procedural Sedation Plan(SUB)*
- Bedside Glucose Nsg
Stat
- IV Insert/Site Care LEB
q2h(std)

Continuous Infusion

- Sodium Chloride 0.9% Bolus
- 20 mL/kg, Injection, IV, once, STAT, (infuse over 15 min), (Bolus) (DEF)**
 - 10 mL/kg, Injection, IV, once, STAT, (infuse over 15 min), (Bolus)*
 - 10 mL/kg, Injection, IV, once, STAT, (infuse over 30 min), (Bolus)*
 - 20 mL/kg, Injection, IV, once, STAT, (infuse over 30 min), (Bolus)*
- Sodium Chloride 0.9%
1,000 mL, IV, STAT, mL/hr
- D5 1/2NS





Physician Orders PEDIATRIC: LEB ED Non-Critical Trauma Plan

- 1,000 mL, IV, STAT, mL/hr
- D5 1/4 NS
1,000 mL, IV, STAT, mL/hr)
- D5 1/2 NS KCl 20 mEq/L
1,000 mL, IV, STAT, mL/hr
- D5 1/4 NS KCl 20 mEq/L
1,000 mL, IV, STAT, mL/hr

Medications

- acetaminophen
 - 15 mg/kg, Liq, PO, once, STAT, Max Dose = 75 mg/kg/day up to 4g/day (DEF)*
 - 325 mg, Tab, PO, once, STAT, Max Dose = 75 mg/kg/day up to 4g/day
- acetaminophen
 - 15 mg/kg, Supp, PR, once, STAT, Max Dose = 75 mg/kg/day up to 4g/day (DEF)*
 - 325 mg, Supp, PR, once, STAT, Max Dose = 75 mg/kg/day up to 4g/day
- acetaminophen-HYDROcodone 325 mg-5 mg oral tablet
1 tab, PO, once, STAT, (1 tab = 5 mg HYDROcodone), Max dose = 10 mg
- acetaminophen-HYDROcodone 325 mg-5 mg oral tablet
2 tab, PO, once, STAT, (1 tab = 5mg of HYDROcodone), Max dose= 10mg
- acetaminophen-HYDROcodone 325 mg-7.5 mg/15 mL oral solution
0.15 mg/kg, Liq, PO, once, STAT, (5mL = 2.5mg HYDROcodone), Max dose= 10mg
- morphine
0.1 mg/kg, Ped Injectable, IV Push, once, STAT, Max initial dose = 10 mg
- morphine
0.1 mg/kg, Ped Injectable, IM, once, STAT, Max initial dose = 10 mg
- HYDROmorphine
0.015 mg/kg, Injection, IV Push, once, STAT, Max initial dose = 0.5 mg
- HYDROmorphine
1 mg, Injection, IV Push, once, STAT
- ondansetron
0.1 mg/kg, Oral Susp, PO, once, STAT, Max dose= 4mg
- ondansetron
4 mg, Orally Disintegrating Tab, PO, once, STAT
- ondansetron
0.1 mg/kg, Ped Injectable, IV Push, once, STAT, Max dose= 8 mg
- LORazepam
0.05 mg/kg, Ped Injectable, IV Push, once, STAT, Max dose: 4 mg





Physician Orders PEDIATRIC: LEB ED Non-Critical Trauma Plan

- EPINEPHrine-lidocaine 1:100,000-1% inj
*mL, Injection, Infiltration, once, STAT, Max dose of lidocaine = 7 mg/kg/dose
Comments: (1 mL = 10 mg of lidocaine component)*
- lidocaine 1% inj
*mL, Injection, Injectable, once, STAT, Max dose of lidocaine = 4.5 mg/kg/dose
Comments: (1 mL = 10 mg)*

Laboratory

- Type and Screen <4 months(DAT included)
STAT, T;N, Reason: to Hold, Type: Blood
- Type and Screen Pediatric
STAT, T;N, Type: Blood
- CBC
STAT, T;N, once, Type: Blood
- BMP
STAT, T;N, once, Type: Blood
- CMP
STAT, T;N, once, Type: Blood
- AST
STAT, T;N, once, Type: Blood
- ALT
STAT, T;N, once, Type: Blood
- BUN
STAT, T;N, once, Type: Blood
- Amylase Level
STAT, T;N, once, Type: Blood
- Amylase Level LeBonheur Germantown
STAT, T;N, once, Type: Blood
- Lipase Level
STAT, T;N, once, Type: Blood
- Lipase Level LeBonheur Germantown
STAT, T;N, once, Type: Blood
- Urinalysis w/Reflex Microscopic Exam
*STAT, T;N, once, Type: Urine, Nurse Collect
If possibility of pregnancy, order one of below:(NOTE)**
- Pregnancy Screen Serum
STAT, T;N, once, Type: Blood

Diagnostic Tests





Physician Orders PEDIATRIC: LEB ED Non-Critical Trauma Plan

- LEB ED Diagnostic X-Ray Plan(SUB)*
- CT Brain/Head WO Cont
T;N, Reason for Exam: Trauma, Stat, Stretcher
- CT Cervical Spine WO Cont
T;N, Reason for Exam: Other, Enter in Comments, Stat, Stretcher
Comments: Trauma
- LEB CT Abdomen W Cont Plan(SUB)*
- LEB CT Chest W Cont Plan(SUB)*
- LEB CT Pelvis W Cont Plan(SUB)*
- CT Maxillofacial Area WO Cont
T;N, Reason for Exam: Trauma, Stat, Stretcher

Consults/Notifications/Referrals

- Consult MD Group
Orthopedic Surgery
- Consult MD Group
General Surgery

Date	Time	Physician's Signature	MD Number
------	------	-----------------------	-----------

***Report Legend:**

- DEF - This order sentence is the default for the selected order
- GOAL - This component is a goal
- IND - This component is an indicator
- INT - This component is an intervention
- IVS - This component is an IV Set
- NOTE - This component is a note
- Rx - This component is a prescription
- SUB - This component is a sub phase, see separate sheet
- R-Required order

